

# Pintoi Peanut: A Seed-propagated Perennial Peanut Forage Option for Florida<sup>1</sup>

Joao M. D. Sanchez, Joao Vendramini, Maria L. Silveira, Jose C. B. Dubeux Jr., Lynn E. Sollenberger, and Philippe Moriel<sup>2</sup>

## IFAS Assessment

Central, North, South

### High Invasion Risk

Predicted to be invasive and not recommended by IFAS. Will be reassessed every 10 years. In particular cases, this species may be considered for use under specific management practices that have been approved by the IFAS Invasive Plant Working Group.

[view assessment](#)

## Introduction

Pintoi peanut (*Arachis pintoii*; also known as pinto peanut, forage peanut, or mani forrajero) is a perennial peanut species native to South America and propagated by seeds. Pintoi peanut and rhizoma peanut (*Arachis glabrata*) are the two most relevant perennial peanut species used as forage; however, the first has been more extensively studied and used in South and Central America, while the second is more common in the southern United States, including Florida. Pintoi peanut is tolerant to grazing and adapted to poorly-drained, low-fertility soils. Because of its adaptability to Florida's environmental conditions and ability to produce viable seeds, pintoi peanut represents an interesting forage alternative for cow-calf producers in the state. This document provides current information on pintoi peanut for forage and livestock producers in Florida.

## Plant Description

Pintoi peanut is a low-growing perennial peanut species that spreads preferentially by stolons (also known as "runners"; Figure 1), without the presence of rhizomes (subterranean stems). The leaflets are different from those of the rhizoma peanut in both color and shape. Both plants have four leaflets in each leaf; however, pintoi peanut leaflets are usually round-tipped and light green, whereas rhizoma peanut leaflets are pointed at the tip and darker green in color (Figure 2). The flowers are predominantly yellow, with some variation between white and reddish orange. The seed production is one of the advantages of pintoi peanut. It occurs in pods, similar

to edible peanut species. At the base of a flower, a peg grows to a length of 2 to 13 inches and subsequently penetrates the soil surface, where it will produce a pod. Pintoi peanut pods usually contain a single seed; however, two or even three seeds may be observed (Figure 3). One of the main characteristics of pintoi peanut pods is that they drop from the pegs after maturity. This makes seed harvest more difficult because it requires soil separation from the pods.



Figure 1. Pintoi peanut stolons (runners) growing parallel to the soil surface.

Credit: J. M. D. Sanchez, UF/IFAS



Figure 2. Comparison between pinto (A) and rhizoma (B) peanut leaflets.

Credit: J. M. D. Sanchez, UF/IFAS



Figure 3. Part of a pinto peanut stolon (runner) with one peg and two pods attached. Like pods of edible peanut, pinto peanut pods remain underground.

Credit: J. M. D. Sanchez, UF/IFAS

## Agronomic Characteristics

Pinto peanut may tolerate acidic (pH = 5.0), low-fertility soils and high soil aluminum concentrations. However, low pH and high soil aluminum concentrations may decrease N fixation and overall pinto peanut production. The plant is adapted to different soil textures ranging from sandy to clay and is more tolerant to poorly drained soils than rhizoma peanut, which makes it an interesting option for producers in south Florida. Recent trials conducted at the UF/IFAS Range Cattle Research and Education Center showed that pinto peanut persisted in soils subjected to intermittent (1- to 2-week) flooding conditions (Sanchez et al. 2018). Reports from International Center for Tropical Agriculture (CIAT) in Colombia indicate that pinto peanut grows well in areas with annual rainfall ranging from 60 to 135 inches, but it also has some drought tolerance (Fisher and Cruz 1994).

Another characteristic that makes pinto peanut a viable forage option in Florida is its frost tolerance. A study conducted in Gainesville, FL found that pinto peanut aboveground biomass was severely damaged by frost, but plant stand survived and new stolons were produced during the rest of the growing season (spring and summer) (Carvalho and Quesenberry 2012). Additionally, pinto peanut can tolerate shade, and it may be used as a cover crop in perennial production systems, such as orange groves.

Previous research in tropical regions showed that dry matter yields of pinto peanut monoculture can range from 100 to 2,500 lb/acre/year, whereas crude protein and in vitro dry matter digestibility concentrations range from 13% to 18%, and from 63% to 67%, respectively (Rincón C. et al. 1992; Carvalho and Quesenberry 2012). The most widely used pinto peanut cultivar is called 'Amarillo'. This cultivar was released in Australia in 1987 and it has satisfactory seed production. The general preference among producers for 'Amarillo' has been based on seed availability. Other genotypes were released in different countries, such as 'Mani Forrajero Perenne' (Colombia in 1992), 'Mani Mejorador' (Costa Rica in 1994), 'Pico Bonito' (Honduras in 1993), 'Belmonte' (Brazil in 1999), 'Amarillo MG-100' (Brazil in 1994), and 'BRS Mandobi' (Brazil in 2011), but seed availability is limited.

## Establishment

Pinto peanut may be established by both stolons and seeds in the spring and summer months. The establishment by seeds requires a seeding rate of 10 lb/acre. Seeding depth in sandy soils should be around 1.5 to 2 inches, whereas seeding in heavier soils should be around 1 inch. The use of specific inoculants may be beneficial for the establishment of pinto peanut, but availability of inoculant in the US may be restricted. The effectiveness of other inoculants, such as commercial products used for cowpea or clover, has not been determined. If vegetative material is used for establishment, stolons should be planted in 4-inch-deep rows, with between-row spacing of 20 inches.

Pinto peanut has slow establishment that may take multiple growing seasons for full ground cover. Research conducted in Gainesville, FL, concluded that pinto peanut ground cover one year after planting was still less than 50% (Carvalho and Quesenberry 2012). There are no registered selective herbicides that can be applied to pinto peanut fields. Herbicides recommended for rhizoma peanut (perennial peanut species commonly used in Florida) may damage pinto peanut stands.

It is recommended to seed pinto peanut with warm-season perennial grasses, such as bahiagrass and bermudagrass. One of the strategies to include pinto peanut in these mixtures is to seed it simultaneously with grass planting. An experiment conducted at the UF/IFAS

Range Cattle Research and Education Center in Ona, FL, showed that seeding pinto peanut either as monoculture or simultaneously with bahiagrass made no difference in pinto peanut spread and productivity (Sanchez et al. 2019). However, when pinto peanut was seeded as monoculture, there was a greater weed infestation even one year after the seeding. Infestation decreased gradually when pinto peanut was seeded in mixture with bahiagrass (Figure 4). Similar observations were reported by Jaramillo et al. (2018b) in a study where pinto peanut was simultaneously planted with 'Tifton-85' bermudagrass (*Cynodon* spp.). Pinto peanut may also be overseeded into existing pastures. It has been observed that pinto peanut established well in preexistent pastures of 'Pensacola' bahiagrass (Jaramillo et al. 2018a) and Jiggs bermudagrass (Sanchez et al. 2018).



Figure 4. Bahiagrass-pinto peanut mixed plot (yellow flowers are from pinto) in broad (A) and close (B) views. Credit: J. M. D. Sanchez, UF/IFAS

## Management

Pinto peanut has a prostrate growth habit with growth points often protected from defoliation at low stubble heights. Researchers in Australia observed that pinto peanut persisted in a grass-legume mixture when stubble heights ranged from 2 to 4 inches (Sinclair et al. 2007). In some pinto peanut-grass mixed pastures, pinto peanut may be favored by lower stubble heights. In Brazil, researchers observed that pinto peanut proportion in a grass-legume mixture with palisadegrass increased from 4% to 14% when stubble height decreased from 14 to 8 inches (Andrade et al. 2012). A few studies were conducted at the UF/IFAS Range Cattle Research and Education Center, evaluating pinto peanut's adaptation to different management practices and its use in mixture with warm-season perennial grasses. Results demonstrated that pinto peanut proportion and spread when overseeded in a Jiggs bermudagrass pasture were similar when grazed at stubble heights of 6 or 12 inches (Sanchez et al. 2018), and that pinto peanut persisted at both stubble heights. Time after planting is an important factor in the initial years after seeding; the legume's proportion in herbage mass increased during the two years of the experiment (Table 1). Another study conducted at the UF/IFAS North Florida Research and Education Center in Marianna, FL, showed that pinto peanut proportion in a grass-legume mixture with Tifton 85 bermudagrass was below 5%

during the first and second years after seeding, reaching 17% on the third year (Jaramillo et al. 2018a). Given its grazing tolerance, low productivity, and prostrate growth habit, pinto peanut should be mainly mixed with other tropical grasses, and used for grazing instead of hay production.

## References

- Andrade, C. M. S. De, R. Garcia, J. F. Valentim, and O. G. Pereira. 2012. "Productivity, Utilization Efficiency and Sward Targets for Mixed Pastures of Marandugrass, Forage Peanut and Tropical Kudzu." *Rev. Bras. Zootec.* 41(3): 512–520. <https://doi.org/10.1590/S1516-35982012000300006>
- Carvalho, M. A., and K. H. Quesenberry. 2012. "Agronomic Evaluation of *Arachis pinto* (Krap. and Gre.) Germplasm in Florida." *Arch. Zootec.* 61(233): 19–29. <https://doi.org/10.4321/S0004-05922012000100003>
- Fisher, M. J., and P. Cruz. 1994. "Some Ecophysiological Characteristics of *Arachis pinto*." In *Biology and Agronomy of Forage Arachis*, edited by P. C. Kerridge and B. Hardy. 169–186. Cali, Colombia: CIAT: Centro Internacional de Agricultura Tropical.
- Jaramillo, D. M., J. C. B. Dubeux, Jr., C. Mackowiak, L. E. Sollenberger, N. DiLorenzo, D. L. Rowland, A. R. S. Blount, E. R. S. Santos, L. Garcia, and M. Ruiz-Moreno. 2018a. "Annual and Perennial Peanut Mixed with 'Pensacola' Bahiagrass in North Florida." *Crop Sci.* 58(2): 982–992. <https://doi.org/10.2135/cropsci2017.09.0542>
- Jaramillo, D. M., J. C. B. Dubeux, Jr., C. Mackowiak, L. E. Sollenberger, E. R. S. Santos, L. Garcia, M. Ruiz-Moreno, C. S. Silva, and N. DiLorenzo. 2018b. "Annual and Perennial Peanut Species as Alternatives to Nitrogen Fertilizer in Bermudagrass Hay Production Systems." *Agron. J.* 110:2390–2399. <https://doi.org/10.2134/agronj2018.01.0036>
- Rincón Castillo, A., P. A. Cuesta Muñoz, R. A. Pérez Bonna, C. E. Lascano, and J. E. Ferguson. 1992. "Maní forrajero perenne (*Arachis pinto* Krapovickas y Gregory): Una alternativa para ganaderos y agricultores." Cali, Colombia.
- Sanchez, J. M. D., J. M. B. Vendramini, M. L. Silveira, L. E. Sollenberger, J. C. B. Dubeux, Jr., P. Moriel, B. Sellers, J. K. Yarborough, and F. C. Leite de Oliveira. 2019. "Seeding Strategies of Bahiagrass and Pinto Peanut Affect Pasture Establishment under Weed Competition." *Grass*

Sanchez, J. M. D., J. M. B. Vendramini, L. E. Sollenberger, M. L. Silveira, J. C. B. Dubeux, Jr., P. Moriel, F. Kuwahara, et al. 2018. "Forage Characteristics of Bermudagrass Pastures Overseeded with Pinto Peanut and Grazed at Different Stubble Heights." *Crop Sci.* 58(4):

Sinclair, K., K. F. Lowe, and K. G. Pembleton. 2007. "Effect of Defoliation Interval and Height on the Growth and Quality of *Arachis pinto* cv. Amarillo." *Trop. Grasslands* 41(4): 260–268.

## Tables

Table 1. Pinto peanut plant population, cover, and biomass production in a grass-legume mixture with Jiggs bermudagrass. Data are expressed according to stubble height treatment (6 or 12 inches), as well as the beginning and end of a two-year experiment (2014 and 2015). Adapted from Sanchez et al. (2018).

Response Variable	Stubble Height (inches) <sup>1</sup>	
	6	12
Pinto peanut ground cover (%)	6	6
Pinto peanut plants/sq. ft.	0.5	0.5
Pinto peanut proportion of total biomass (%)	3	3

  

Response Variable	Year <sup>2</sup>	
	2014 (Initial)	2015 (Final)
Pinto peanut proportion of total biomass (%)	1	8

<sup>1</sup> Data refer to the average of two experimental years.

<sup>2</sup> Data refer to the average of 6- and 12-inch stubble height treatments.

<sup>1</sup> This document is SS-AGR-445, one of a series of the Agronomy Department, UF/IFAS Extension. Original publication date November 2020. Visit the Ask IFAS website at <https://ask.ifas.ufl.edu> for the currently supported version of this publication.

<sup>2</sup> Joao M. D. Sanchez, former research assistant (Ph. D. 2018, Agronomy), UF/IFAS Range Cattle Research and Education Center, Ona, FL; João Vendramini, center director, Texas A&M AgriLife Stephenville Research and Extension Center, former professor and forage specialist, Department of Agronomy, UF/IFAS Range Cattle Research and Education Center; Maria L. Silveira, former professor, soil fertility and water quality, Department of Soil, Water, and Ecosystem Sciences, UF/IFAS Range Cattle Research and Education Center, Ona, FL; José C. Dubeux, professor of forage management, Department of Agronomy, UF/IFAS North Florida Research and Education Center; Lynn E. Sollenberger, distinguished professor of forage crops, UF/IFAS Agronomy Department, Gainesville, FL; Philippe Moriel, associate professor, Department of Animal Sciences, UF/IFAS Range Cattle Research and Education Center; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information, and other services only to individuals and institutions that operate in compliance with applicable federal and state non-discrimination laws and policies. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A&M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, Dean for UF/IFAS Extension.