

2026–2029 Florida Citrus Production Guide: Citrus Root Weevils¹

Larry W. Duncan, Lukasz Stelinski, and Lauren M. Diepenbrock²

Citrus root weevils represent a complex of species known to infest citrus trees and various alternate host plants in Florida. The most common species infesting citrus, in order of greatest geographical distribution, are the *Diaprepes* root weevil, *Diaprepes abbreviatus*; the blue-green citrus root weevils, *Pachnaeus litus* and *Pachnaeus opalus*; the little leaf notcher, *Artipus floridanus*; and the Fuller rose beetle, *Asynonychus godmani*. Other lesser species inhabit citrus on occasion.

All citrus root weevils have a similar life cycle. They have three immature stages: egg, larva, and pupa. Adult weevils emerge from the soil and lay eggs on host plants aboveground, the larvae drop to the soil to feed on roots, and the pupae and teneral adult stages are spent belowground. Adults emerge from the soil throughout the year. Peak emergence varies within species and by geographical region (ridge vs. coastal and interior flatwoods). Peak adult emergence for the blue-green root weevils and Fuller rose beetle is normally April and May. *Diaprepes* adult emergence from the soil peaks in late May to early July, while peak adult abundance on the tree canopy parallels adult emergence in May/June but can have a second peak in late August to mid-October. The second peak is sporadic. The little leaf notcher has three generations per year. Although there is some overlap of generations, adults appear most abundant on trees in April/May, July/August, and October/November. All adult weevils are attracted to the nonreflective silhouette of the citrus tree trunk. The little leaf notcher and Fuller rose beetle are flightless and must crawl up the trunk, but other species will fly to the canopy.

The most visible plant damage resulting from adult feeding is notching of the margins of leaves of young, tender shoots. Notching patterns differ slightly among species and can be confused with grasshopper injury. Prolonged leaf feeding by adults appears to cause no economic effects in mature groves; however, on occasion, feeding will cause virtual defoliation of small replants.

With the exception of little leaf notchers, which prefer a weed host, larval feeding injury to the roots by other root weevils, particularly *Diaprepes* root weevil, can have a devastating effect on citrus trees because all larval stages

feed on the roots for most of the year. Tiny hatchlings feed on fibrous roots, whereas larger larvae feed on the larger structural roots, forming deep grooves as they consume the outer bark, including the cambium layer. Roots may be girdled and killed in the process, or the crown may be girdled, causing tree death. Larval feeding sites predispose the root system to infection and girdling by *Phytophthora* spp., thereby exacerbating economic loss.

Pest Management

Methods of Sampling Root Weevil Larvae and Adults

The population abundance and distribution of endemic citrus root weevils, regardless of species, vary from grove to grove, within a grove, and within a season. The peak seasonal soil emergence of adults within a citrus grove can be monitored using ground traps to capture emerging adults or via visual sighting of adults in the tree canopy. No methods exist for monitoring larvae in the soil. By monitoring adult emergence and abundance, the approximate time and intensity of peak adult emergence can be estimated for each infestation. Depending on the species of weevil and their seasonal abundance pattern, a grower can apply adult control measures when weevil populations are highest. Research suggests that adult emergence in spring often coincides with the onset of summer rains in late May through June, increasing soil temperature and the summer flush in central Florida groves.

Cultural Considerations

Citrus root weevil management begins with the selection of a phytophthora-resistant rootstock that is certified weevil-free. Optimal soil drainage is fundamental to citrus root weevil management, particularly in heavier soils common to the coastal and interior flatwoods where insect and pathogen populations are highest. Tree decline associated with *Diaprepes* distribution is often patchy within groves and most obvious in lowlands. Stressed trees frequently harbor higher populations of adults because these stressed trees frequently generate more leaf flushes as food for adults. Spot-treating these locations with a chemical or biological agent should help. Regular

fertilization and irrigation are crucial to new root growth in weevil-infested groves. Fertigation at monthly intervals has been used by growers to promote the growth of fibrous roots after *Diaprepes* has destroyed the taproot and inner crown of the tree. Skirt pruning and trunk banding can be effective in controlling flightless weevil species. Weed control is also needed to prevent movement into trees from stems of grasses and/or broadleaf weeds. Weed control is probably beneficial in reducing populations of alternate host plants. **The use of sound cultural practices by the grower should be adequate for managing all citrus root weevils on mature trees except for the *Diaprepes* root weevil and blue-green citrus root weevils.**

A wide range of parasites, predators, and pathogens attack citrus root weevils at one or more developmental stages within the tree canopy or in the soil. Most of these natural enemies are widely distributed and are general feeders. When focusing on cultural tactics favoring tree health and not using chemical methods, growers are conserving and augmenting the natural enemies of citrus root weevils.

Pest Control Considerations

Pest management of *Diaprepes* and, to a lesser extent, other citrus root weevils must begin with control of different life stages, particularly adult weevils, using the following options: (1) foliar sprays for egg and adult suppression, (2) chemical barriers for larval control, and (3) biological control of all subterranean stages with nematodes. The application of these control tactics is timed according to monitoring of adult emergence and the onset of leaf flushing in the spring/summer period. Any of these tactics should reduce root injury and help sustain root health from grove to grove. For many groves, however, pest management might differ according to (1) rootstock susceptibility to soilborne diseases (i.e., *Phytophthora* spp.) and (2) root stress caused by excessive flooding and poor drainage of sandy loam soils. In certain grove situations, a soil fungicide for control of *Phytophthora* spp. should be advised.

Newly planted resets and groves younger than five years old with an established *Diaprepes* infestation on a susceptible rootstock can decline within two years without adult and/or larval control. A similar grove situation involving a resistant rootstock will have lesser tree decline but will require adult suppression. **Remember, groves planted on deep, sandy soils will often require no supplemental control and can rely on biological control agents.**

Foliar sprays of different contact (knockdown) insecticides that include petroleum oil to improve residual effect are used to target adult weevils in the tree canopy. Although foliar sprays have been used by growers to suppress adults any time of the year, research in central Florida has shown conclusively that root injury is lessened

and overall tree health improved when two foliar sprays are used 4 weeks apart during peak summer flush in late May through June, along with an egg sterilant in the last application. The purpose of adult suppression with foliar sprays is to limit the number of gravid females and egg deposition, thereby reducing the number of larvae entering the soil. An egg sterilant such as Micromite 80WGS has a six-week residual effect, during which females lay sterile eggs and eggs contacting the leaf surface are nonviable. The addition of petroleum oil to the spray mixture affects the bonding characteristics of the substance bonding the egg mass to the leaf.

Multiple applications of most foliar sprays within a season can incite an abnormal increase in spider mite populations; any pesticide, when used frequently, might cause secondary pest outbreaks or lead to resistance.

A chemical barrier applied as a band to the soil surface beneath the tree through an herbicide applicator provides a treated surface that will kill newly hatched invasive larvae before they reach the root system. The chemical must be uniformly applied from the trunk to the dripline of the tree to a moistened soil surface devoid of litter. Greater spray volume (~40 gal/ac) should ensure greater uniformity of coverage. Disturbance of the soil beneath the trees should be minimized to protect the soil barrier. Because neonates are killed upon exposure to treated soil as they pass through the barrier, this control tactic is best used for resets and young plantings infested with phytophthora, where root injury by larvae must be minimal.

Timing chemical application to the time of year when larval entry into the soil is highest requires monitoring of adult weevils in the tree. Because highest larva recruitment occurs just after peak adult emergence, growers should apply soil treatment in early July, about two weeks after peak adult emergence in central Florida. Peak adult emergence is generally two to three weeks earlier in coastal groves.

Currently, Brigade WSB, a synthetic pyrethroid, is the only chemical registered for neonatal larvae control and applied as a soil barrier. Brigade has about three weeks of residual presence in the soil and will suppress ants foraging on the soil surface. Generally, ant predators will recover after 30 days.

Physical barriers consisting of landscape fabric on the soil surface were shown experimentally to be highly effective in preventing the newly hatched larvae from entering the soil and newly developed adults from escaping beneath the fabric. These barriers have been used for weevil management in California and Texas with some success and are increasingly employed in Florida on both flat-ground and raised beds. In addition to weevil management, the fabric has been shown to conserve water

use and increase root and tree growth rates. Various ground-cover fabric products have been used in citrus trials (e.g., Lumite, Ultraweb 3000), but it is critical to use a product with high water infiltration rate to avoid irrigation runoff. The cost of the fabric and installation is high, but the savings in reduced water use and weed and insect pest management, as well as better tree growth and reduced weevil damage, would potentially make the treatment profitable where weevil damage is high.

The use of individual protective tree covers (IPC) for psyllid management will also prevent adult *Diaprepes* from laying eggs on young trees. Protecting trees from both huanglongbing and *Diaprepes* root feeding during the first several years, when trees are most susceptible to damage, makes the use of these devices an especially promising IPM tactic.

Parasitic nematodes that specifically attack insects are infectious to all larval stages of citrus root weevils. They are naturally found in citrus soils, where they inflict mortality to all weevil life stages they contact. Depending on availability, nematodes are also sold as biopesticides to control citrus root weevil larvae. They should be applied during months when soil surface temperatures are expected to exceed 70°F. Weevil larvae are generally most abundant in the soil during the summer (mid-July through September); therefore, one or more nematode applications

are recommended at this time of year if soil moisture via natural causes and/or irrigation is adequate. Nematodes should not be applied within four weeks of nematicide use. Properly modified herbicide applicators or microsprinkler irrigation systems are used to deliver nematodes into premoistened soil. Approximately one acre-inch of water should also be applied to the irrigated acre immediately following application. Application late in the day or on cloudy days is encouraged to reduce nematode desiccation and exposure to lethal UV radiation.

Nematode products are most effective when applied in sandy soils with coarser soil texture and are less effective in very fine-textured soils at recommended rates. Higher rates can be applied to very fine-textured soils. *Nematode products that have been tested and recommended by UF/IFAS are currently unavailable.*

A fungicide for control of *Phytophthora* spp. may be recommended under the following conditions as a supplemental strategy to larval and adult weevil control: (1) the soils are fine-textured, poorly drained, or high in pH and calcium carbonate; (2) the trees are on rootstocks susceptible to *Phytophthora* spp.; and (3) populations are above the damaging levels (20 and 40 propagules per cm³ soil) for *P. nicotianae* and *P. palmivora*, respectively. Remember, larval and/or adult weevil control must be effective before fungicide treatment is justified.

¹ This document is ENY-611, one of a series of the Department of Entomology and Nematology, UF/IFAS Extension. Original publication date July 2000. Revised annually. Most recent revision August 2026. Visit the Ask IFAS website at <https://ask.ifas.ufl.edu> for the currently supported version of this publication. © 2026 UF/IFAS. This publication is licensed under [CC BY-NC-ND 4.0](#).

² Larry W. Duncan, professor, nematology, Department of Entomology and Nematology, UF/IFAS Citrus Research and Education Center; Lukasz Stelinski, professor, Department of Entomology and Nematology, UF/IFAS Citrus Research and Education Center; Lauren M. Diepenbrock, associate professor, integrated pest management and insect ecology, Department of Entomology and Nematology, UF/IFAS Citrus Research and Education Center, Lake Alfred, FL; UF/IFAS Extension, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition.

Use pesticides safely. Read and follow directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information, and other services only to individuals and institutions that operate in compliance with applicable federal and state non-discrimination laws and policies. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A&M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, Dean for UF/IFAS Extension.