

2026–2029 Florida Citrus Production Guide: Pesticide Resistance and Resistance Management¹

Lauren M. Diepenbrock, Megan M. Dewdney, and Ramdas Kanissery²

Populations of animals, fungi, bacteria, and plants possess the ability to respond to sustained changes or stresses in their environment in ways that enable the continued survival of the species. Such environmental stresses include physical factors (e.g., temperature or humidity), biological factors (e.g., predators, parasites, or pathogens), and environmental contaminants. In any population, a small percentage of individuals will be better able to respond to new stresses because of unique traits or characteristics that they possess. Consequently, those individuals will survive, reproduce, and become more common in a population. This phenomenon is commonly referred to as "survival of the fittest."

Many pest species, such as the citrus rust mite, are exceptionally well equipped to respond to environmental stresses because of their short generation time and large reproductive potential. The use of chemical sprays to control insects, mites, bacterial and fungal diseases, and weeds of citrus creates a potent environmental stress. There are now many examples of pests, pathogens, and weeds that have responded by developing resistance to one or more pesticides. Pesticide-resistant individuals are those that have developed the ability to tolerate doses of a toxicant that would be lethal to the majority of individuals. The resistance mechanisms can vary according to pest species and/or the class of chemical to which the pest is exposed. Resistance mechanisms for insects include an increased capacity to detoxify the pesticide once it has entered the pest's body, a decreased sensitivity of the target site that the pesticide acts upon, a decreased penetration of the pesticide through the cuticle, or sequestration of the pesticide within the organism. The main resistance mechanism for fungal pathogens is a change in the target site so that the pathogen is less susceptible or fully resistant. With repeated or intense exposure to herbicides, some weeds develop resistance because only individuals that are capable of detoxifying the chemical persist over time. A single resistance mechanism can sometimes provide defense against different classes of chemicals; this is known as *cross-resistance*. When more than one resistance mechanism is expressed in the same individual, this individual is said to show *multiple resistance*.

Of the factors that affect the development of resistance, including the pest's or pathogen's biology, ecology, and genetics, only the operational factors can be manipulated by the grower. The key operational factor that will delay the onset of pesticidal resistance and prolong the effective life of a compound is assuring the survival of some susceptible individuals to dilute the population of resistant individuals. The following operational procedures should be on a grower's checklist to steward sound pesticidal resistance management for acaricides, insecticides, fungicides, antibiotics, and herbicides:

1. Never rely on a single pesticide class.
2. Integrate chemical control with effective and complementary cultural and biological control practices.
3. Always use pesticides at recommended rates and strive for thorough coverage.
4. When there is more than one generation of pest, alternate different pesticide classes.
5. Do not use tank mixtures of products that have the same mode of action.
6. If control with a pesticide fails, do not re-treat with a chemical that has the same mode of action.

Reports of resistance have been documented for certain acaricides used to control citrus rust mite and fungicides used to combat diseases in Florida. Resistance is also likely to be found in weeds with repeated exposure to certain herbicides. Resistance to Benlate developed in the greasy spot fungus shortly after the product was introduced about 30 years ago and is still widespread. More recently, we have discovered that there is QoI fungicide (Table 2) resistance in the greasy spot fungus. This is a developing story, but caution should be used when choosing this class of fungicide, and a mixture of modes of action is advised. Benlate resistance also occurs in the scab fungus in isolated situations and is stable. In tangerine groves with Alternaria brown spot, resistance has been detected to strobilurin fungicides (Gem, Headline, and Quadris [formerly Abound] and contained in the mixtures Pristine, Priaxor, and Amistar Top), but no resistance has developed to ferbam. Unfortunately, ferbam was deregistered in 2025, removing a valuable resistance management tool.

Dicofol resistance in citrus rust mite was detected throughout the citrus industry about 10 years ago, but resistance proved to be unstable, and usage of dicofol has continued. Agri-mek tolerance in citrus rust mite is of concern, and growers should follow sound resistance management practices when using this product. Recent studies have shown reduced susceptibility to several insecticides in populations of Asian citrus psyllid after repeated exposure to similar materials, but that susceptibility can be restored by rotating modes of action used in management programs. Resistance management is crucial to the management of this insect. Glyphosate-resistant weeds are becoming commonplace in many production systems with the repeated use of this popular preemergence herbicide, highlighting the need to rotate materials for weed management.

The following tables are provided to aid in the rotation of pesticides with different modes of action within a season or from year to year. There are separate tables for insecticides/acaricides, fungicides/antibiotics, and herbicides. The information in these tables was derived from information produced by the [Insecticide Resistance Action Committee \(IRAC\)](#), the [Fungicide Resistance Action Committee \(FRAC\)](#), and the [Herbicide Resistance Action Committee \(HRAC\)](#). Each table lists the number (or letter in the case of herbicides) of the group code for each pesticide class, the group name or general description of that group of pesticides, the common name of pesticides used in citrus production that belong to each group, and examples of trade names of pesticides for each common name listed. When using the table to rotate between using products with different modes of action, choose products with a different group code than previously used in the grove during the current growing season. In the case of insecticides/acaricides, many of these pesticides are broken into subgroups. It is unclear whether cross-resistance will occur between these subgroups. When possible, it is recommended to rotate with an entirely different group. (Note: The IRAC and FRAC mode of action systems both use a similar numbering system. There is no cross-resistance potential between the insecticides and fungicides.) Products with broad-based activity such as sulfur and oil are not included in this list because the development of resistance to them is not likely.

Web Addresses for Links

FRAC: <https://www.frac.info/>

HRAC: <https://hracglobal.com/tools/classification-lookup>

IRAC: <http://www.irac-online.org/>

Table 1. Insecticides and miticides used in Florida citrus grouped by mode of action.

IRAC Group ¹	Subgroup	Group Name	Common Name	Trade Name
1	1A	Carbamates	carbaryl oxamyl	Sevin Vydate, ReTurn
	1B	Organophosphates	acephate dimethoate malathion naled phosmet	Orthene, Acenthrin, Avatar PLX, Bracket Dimethoate Malathion Dibrom Imidan
3	3A	Pyrethroids	bifenthrin fenpropathrin zeta-cypermethrin	Brigade Danitol Mustang
4	4A	Neonicotinoids	acetamiprid clothianidin imidacloprid thiamethoxam	Actara, Admire Pro, Advise, Alias, Assail, Belay, Couraze, Imida E-Ag, Impulse, Macho, Montana, Nuprid, Pasada, Platinum, Prey, Torrent, Widow
	4D	Butenolides	flupyradifurone	Sivanto, Altus
5		Spinosyns	spinosad spinetoram	Spintor Delegate
6		Avermectins	abamectin	Abacus, Abba, Agri-mek, Clinch, Epi-mek, Reaper, Zoro
7	7A	Juvenile Hormone Analogues	methoprene	Extinguish Ant Bait
	7B	Fenoxycarb	fenoxycarb	Precision
	7C	Pyriproxyfen	pyriproxyfen	Knack
10	10A	Hexythiazox	hexythiazox	Savey
11	11A	<i>Bacillus thuringiensis</i> (Bt)	<i>Bt</i> var. aizawai	Various
			<i>Bt</i> var. kurstaki	Various
12	12B	Organotin miticides	fenbutatin oxide	Vendex
	12C	Propargite	propargite	Comite
15		Benzoylureas	diflubenzuron	Micromite
16		Buprofezin	buprofezin	Applaud, Centaur
18		Diacylhydrazines	methoxyfenozide	Intrepid
21	21A	METI acaricides	pyridaben	Nexter
			fenpyroximate	Portal, Apta
23		Tetronic/Tetramic acid derivatives	spirodiclofen	Envidor
			spirotetramat	Movento, Senstar
28		Diamides	chlorantraniliprole, cyantraniliprole	Exirel, Verimark, Voliam Flexi (one component)

IRAC Group ¹	Subgroup	Group Name	Common Name	Trade Name
30		Meta-diamides & Isoxazolines	plinazolin	Zivalgo
UN		Unknown MOA	bifenazate	Acramite
			cryolite	Kryocide
			dicofol	Kelthane
¹ Mode of action based on the Insecticide Resistance Action Committee (IRAC) Mode of Action Classification v. 11.4 (2025).				

Table 2. Fungicides and antibiotics used in Florida citrus grouped by mode of action.

FRAC Group ¹	Group Name	Common Name	Trade Name
1	MBC—fungicides (Methyl benzimidazole carbamates)	thiabendazole	Many (TBZ)
3	DMI—fungicides (Demethylation inhibitors)	difenoconazole fenbuconazole imazalil mefentrifluconazole propiconazole	Amistar Top, Miravis Top, Acadia ESQ, Luna Flex, Tycoon, and others Enable 2F, Indar 2F Many Cevya, Provysol, Veltyma Many
4	PA—fungicides (Phenylamides)	metalaxyl mefenoxam	Ridomil, ReCon, Regulate, MetaStar, Xyler Ultra Flourish, Ridomil Gold, Subdue, Mefenoxam 2 AQ, ReCon Bold, Stergo MX, others
7	SDHI—fungicides (Succinate-dehydrogenase inhibitors)	boscalid fluopyram fluxapyroxad pydiflumetofen	Pristine, Endura, Bonafide, Empire Luna Sensation, Luna Flex, Broadform, Velum Priaxor Xemium, Republic, Tesaris Miravis Top
11	QoI—fungicides (Quinone outside inhibitors)	azoxystrobin trifloxystrobin pyraclostrobin	Quadris (formerly Abound) and others, Graduate A+, Amistar Top Gem, Broadform, Compass, Luna Sensation, Venus Headline and others, Pristine, Priaxor Xemium, Veltyma
12	PP—fungicides (Phenylpyrroles)	fludioxonil	Graduate, Graduate A+, Pilato SC
19	Polyoxin	polyoxin D zinc salt	OSO 5% SC, Ph-D
40	CAA—fungicide (Carboxylic acid amides)	mandipropamid	Revus Orondis Ultra
41	Tetracycline antibiotic	oxytetracycline	Fireline, Mycosheild ReMedium TI, Rectify
43	Benzamides	fluopicolide	Adorn, Presidio

FRAC Group ¹	Group Name	Common Name	Trade Name
48	Polyene	natamycin	BioSpectra 100SC
49	OSBPI—oxysterol binding protein homologue inhibition	oxathiapiprolin	Orondis Orondis Ultra
M 01	Inorganic	copper	Many
P 01	Benzothiadiazole (BTH)	acibenzolar-S-methyl	Actigard 50WG
P 07	Phosphonates	fosetyl-Al phosphorous acid and salts	Aliette, Linebacker Many
¹ Mode of action based on the 2025 Fungicide Resistance Action Committee (FRAC) Mode of Action Classification.			

Table 3. Herbicides used in Florida citrus grouped by mode of action.

HRAC Group ¹	Group Name/Chemical Family	Common Name	Trade Name(s)
A (1)	Aryloxyphenoxy-propionates Cyclohexanediones	fluazifop-p-butyl sethoxydim	Fusilade Poast
C1 (5)	Triazines Uracils	simazine bromacil	Caliber, Princep, Simazine Hyvar, Krovar
C2 (5)	Ureas	diuron	Direx, Karmex, Krovar
D (22)	Pyridiniums	paraquat	Gramoxone
E (14)	N-Phenyl-triazolinones N-Phenyl-imides	carfentrazone-ethyl flumioxazin saflufenacil	Aim Chateau EZ Treevix
F1 (12)	N-Phenyl heterocycles	norflurazon	Solicam
F2 (27)	Triketones	mesotrione	Broadworks
G (9)	Glycine	glyphosate	(Various) e.g., Roundup, Gly Star
H (10)	Phosphinic acids	glufosinate-ammonium	Rely 280, Scout
K1 (3)	Dinitroanilines	oryzalin pendimethalin	Surflan Prowl
L (29)	Alkylazines	indaziflam	Alion
O (4)	Phenoxy-carboxylates	2,4-D	Embed Extra
¹ Mode of action based on the 2025 Herbicide Resistance Action Committee (HRAC) Mode of Action Classification. HRAC/Weed Science Society of America (WSSA) numerical codes for this classification are provided in parentheses.			

¹ This document is ENY-624, one of a series of the Department of Entomology and Nematology, UF/IFAS Extension. Original publication date December 1995. Revised annually. Most recent revision August 2026. Visit the Ask IFAS website at <https://ask.ifas.ufl.edu> for the currently supported version of this publication. © 2026 UF/IFAS. This publication is licensed under [CC BY-NC-ND 4.0](#).

² Lauren M. Diepenbrock, associate professor, integrated pest management and insect ecology, Department of Entomology and Nematology, UF/IFAS Citrus Research and Education Center, Lake Alfred, FL; Megan M. Dewdney, professor and Extension specialist, Department of Plant Pathology, UF/IFAS Citrus Research and Education Center, Lake Alfred, FL; Ramdas Kaniserry, associate professor, weed science, Department of Horticultural Sciences, UF/IFAS Southwest Florida Research and Education Center, Immokalee, FL; UF/IFAS Extension, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition.

Use pesticides safely. Read and follow directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information, and other services only to individuals and institutions that operate in compliance with applicable federal and state non-discrimination laws and policies. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A&M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, Dean for UF/IFAS Extension.