

Strawberry Star™ Caladium—A Spotted, Fancy-Leaved Variety for Use in Containers and Landscapes¹

Zhanao Deng²

Caladiums are grown in containers and landscapes for their bright, colorful leaves. Commercially available caladium plants are generally produced from tubers. Some existing commercial caladium varieties have attractive colors or coloration patterns but poor container and landscape performance and/or poor tuber yield. 'Marie Moir' is an example of this type of variety (Wilfret and Hurner 1982). Its leaves are very attractive but are few, resulting in poor container and landscape displays that are unsatisfactory for commercial or homeowner acceptance. Additionally, its field production of tubers has been unstable over the years, and caladium growers cannot profitably grow the variety.

Strawberry Star™ caladium is a fancy-leaved variety with a primarily white leaf face and attractive red spots (Figure 1) (Deng and Harbaugh 2011). It is similar to 'Marie Moir' in leaf color and coloration pattern but different from 'Marie Moir' in petiole color (green vs. brown on 'Marie Moir'). Compared to 'Marie Moir', Strawberry Star™ caladium sprouts earlier, produces high-quality container plants, demonstrates better landscape performance, and yields more tubers. These improvements make Strawberry Star™ caladium a desirable replacement for 'Marie Moir' and an additional sun-tolerant variety for the landscape plant palette.



Figure 1. Strawberry Star™ caladium plants grown in ground beds in full sun

Credit: Zhanao Deng, University of Florida

Strawberry Star™ caladium is distinct from Cranberry Star™ caladium, a variety released in 2007 that has burgundy/purple spots on its leaves. Strawberry Star™ caladium is much improved over Cranberry Star™ caladium in sunburn tolerance and landscape performance.

Origin

Strawberry Star™ caladium resulted from a cross between 'Summer Rose' and 'Florida Fantasy' and was released under the name 'UF-85-5'. 'Summer Rose' was the progeny of 'Aaron' and University of Florida caladium breeding line UF-FCT, which resulted from a cross between 'Fire Chief' and 'Torchy'. 'Florida Fantasy' was the progeny of 'Candidum Junior' and 'Red Frill'. The ancestry of 'Aaron', 'Candidum Junior', 'Fire Chief', 'Red Frill', and 'Torchy' is unknown.

Leaf and Tuber Characteristics

Leaves of Strawberry Star™ caladium are heart shaped and have palmate-pinnate venation. The upper leaf surface has a green margin bordering the entire leaf, several green main veins, and numerous secondary green veins, which are netted over the whole leaf blade (Figure 1). Interveinal areas are mostly green-white. Red spots occur between main veins in the range of 10–30 per fully developed leaf. The undersurface has a yellow-green undertone with heavily netted veins of gray-green. Primary veins are gray-green at the basal notch and near the tip of the leaf. Spots are gray-purple and vary in size. The prominent petiole color is yellow-green with streaks of gray-brown to brown. Jumbo-sized (1.5–2.5 inches in diameter) tubers have multiple segments and bear four to six dominant buds. The tuber surface is brown, and the cortical area is yellow.

Tuber Production

Strawberry Star™ caladium was evaluated for tuber production under the field conditions at the Gulf Coast Research and Education Center in Wimauma, Florida, in 2005 and 2006. The soil was EauGallie fine sand with about 1% organic matter and a pH between 6.2 and 7.4. Plants were grown in a plastic-mulched, raised-bed system with a constant water table maintained through seepage irrigation. On February 25, 2005, ground beds

were fumigated with a mixture of 67% methyl bromide and 33% chloropicrin (by volume) at the rate of 350 pounds per acre. The raised beds were 32 inches wide and 8 inches high. Caladium seed pieces (cut tuber propagules ~1 inch x ~1 inch x ~1 inch) were planted in early to mid-April with 6 inches in-row and between-row spacing. Osmocote®, a controlled-release fertilizer (18-9-12, 8–9 months), was manually applied to the bed surface at the rate of 300 pounds of nitrogen (N) per acre when caladium shoot tips were emerging from the soil. Tubers (new crop) were harvested in late November 2005. Harvested tubers were washed, dried, weighed, counted, and graded. Tuber grading was by maximum diameter: No. 2 (1–1.5 inches), No. 1 (1.5–2.5 inches), Jumbo (2.5–3.5 inches), Mammoth (3.5–4.5 inches), and Super Mammoth (> 4.5 inches). Tuber counts and grades were converted into a production index (PI) to show the relative economic value of the harvested tubers per experimental plot: $PI = n(\text{No. 2}) + 2n(\text{No. 1}) + 4n(\text{Jumbo}) + 6n(\text{Mammoth}) + 8n(\text{Super Mammoth})$, where n is the number of tubers in each grade. The same cultural practices were used in 2006, except that the beds were fumigated on March 10, 2006, with the same fumigant mixture but at the rate of 175 pounds per acre, and tubers were harvested in mid-December 2006.

In both 2005 and 2006, 13-square-foot field plots were organized in three randomized complete blocks, each with 30 plants. Three commercial varieties, 'Candidum', 'Marie Moir', and 'Miss Muffet', were included in the experiment for comparison. 'Candidum' is the No. 1 or No. 2 best-selling caladium variety (Bell, Wilfret, and DeVoll 1998; Deng et al. 2008) and shares a similar coloration pattern (without the spots) with Strawberry Star™ caladium. As mentioned previously, 'Marie Moir' is the closest to Strawberry Star™ caladium in leaf coloration among commercial varieties. 'Miss Muffet' was included because it is a popular spotted variety.

Strawberry Star™ caladium was significantly more productive than 'Candidum' and 'Miss Muffet' in both the 2005 and 2006 growing seasons (Table 1). Its average tuber weight was 115% (2005) and 83% (2006) greater than that of 'Candidum' and 75% (2005) and 103% (2006) greater than that of 'Miss Muffet'. Its production index was 66% (2005) and 53% (2006) higher than that of 'Candidum' and 49% (2005) and 65% (2006) higher than that of 'Miss Muffet'. Its number of marketable tubers was 50% (2005) and 44% (2006) larger than that of 'Candidum' and 36% (2005) and 28% (2006) larger than that of 'Miss Muffet'. Strawberry Star™ caladium showed good consistency in tuber weight, number of marketable tubers, and production index between growing seasons, while 'Marie Moir' had poor consistency between growing seasons in tuber production. 'Marie Moir' and Strawberry Star™ caladium yielded similarly well in 2005, but 'Marie Moir' yielded poorly in 2006—approximately 33% less than the tuber weight and production index of Strawberry Star™

caladium and only about 50% of the number of marketable tubers for Strawberry Star™ caladium. These results are similar to growers' reports of 'Marie Moir' over the years; that is, 'Marie Moir' is not a dependable variety from year to year.

Container Forcing and Performance

The suitability of Strawberry Star™ caladium for container forcing was evaluated by forcing tubers in 4.5-inch containers. No. 1 tubers of Strawberry Star™ caladium, 'Marie Moir', and 'Miss Muffet' were planted either intact or de-eyed in a peat/vermiculite mix (VerGro Container Mix A, Verlite, Tampa, FL) in March 2007. Container plants were grown on metal benches in a greenhouse with 45% light exclusion. Temperatures in the greenhouse ranged from 61°F (night) to 85°F (day) during the experiment.

Tubers of Strawberry Star™ caladium sprouted in 30 days (intact) or 32 days (de-eyed) after planting, similar to 'Miss Muffet' plants, but 4–9 days earlier than 'Marie Moir' plants (Table 2). Container-grown Strawberry Star™ caladium plants were 7–9 inches tall, regardless of tuber treatment (intact or de-eyed), similar to plants of 'Marie Moir' but significantly taller (2–3 inches) than 'Miss Muffet' plants. Strawberry Star™ caladium produced an average of 12 leaves per intact plant 8 weeks after planting and 21 per de-eyed plant, similar to 'Miss Muffet' but significantly more than 'Marie Moir'. Leaves of Strawberry Star™ caladium were significantly smaller than those of 'Marie Moir' when tubers were planted intact (~3 inches shorter and ~1 inch narrower). When tubers were de-eyed, leaf size differences became smaller (~0.8 inches shorter and 0.8 inches narrower). With numerous medium-sized leaves, Strawberry Star™ caladium container plants were of significantly higher quality (3.4–4.4) than those of 'Marie Moir' (1.9–2.4). Strawberry Star™ caladium container plant quality ratings were similar to those of 'Miss Muffet', which is known for its good growth habit. Container trials indicated that tuber de-eyeing results in higher plant quality ratings for Strawberry Star™ caladium (Table 2).

Landscape Performance

The landscape performance of Strawberry Star™ caladium under full-sun conditions was evaluated in 2005 and 2006 in the same field plots used for evaluating tuber production. 'Candidum', 'Marie Moir', and 'Miss Muffet' were included in the tests for comparison. The overall plant performance was rated multiple times (June, July, and August) in each growing season on a scale of 1–5, with 1 being very poor (few leaves and lack of vigor) and 5 being excellent (full plants, numerous leaves, and bright color display). Leaf sunburn tolerance was also evaluated multiple times in each growing season on a scale of 1–5, with 1 being very susceptible to sunburn and showing numerous sun-damaged areas or holes on leaves and 5 being resistant

to sunburn and not showing any sun-damaged areas. At approximately 4 months after planting, plant height, number of leaves, and leaf sizes were measured.

Strawberry Star™ caladium plants were 4.1–5.4 inches taller and developed significantly more (~100%) leaves than 'Candidum' and 'Marie Moir' (Table 3). Leaf sizes were similar among Strawberry Star™ caladium, 'Candidum', and 'Marie Moir' (9.4–9.8 inches long and 5.9–6.1 inches wide). 'Miss Muffet', a dwarf variety, had shorter plants (7.1 inches) and smaller leaves (on average, 6.6 inches long and 4.3 inches wide).

Strawberry Star™ caladium plants performed well in the landscape, with performance ratings between 3.6 and 4.8, significantly higher than those of 'Candidum' (1.7–3.1), 'Marie Moir' (1.4–3.3), and 'Miss Muffet' (1.5–2.4). Strawberry Star™ caladium received the highest sunburn tolerance ratings in all the evaluations in each growing season, indicating better sun tolerance in this variety compared to the commercial varieties and potential for use in sunny locations in the landscape. Strawberry Star™ caladium showed a high level of similarity to Cranberry Star™ caladium in plant height, leaf number, and leaf size, but Cranberry Star™ caladium is sensitive to full sun (Table 3) and thus is limited to use in shady locations in the landscape.

Recommendation

Strawberry Star™ caladium sprouts 4–9 days earlier, produces higher-quality container plants, and performs better in the landscape than 'Marie Moir', and it has shown better sunburn tolerance than Cranberry Star™ caladium. Additionally, Strawberry Star™ caladium has shown improved tuber yield potential over 'Marie Moir', 'Candidum', and 'Miss Muffet'. In container plant production, Strawberry Star™ caladium behaves like 'Miss Muffet', with a similar number of days to sprout, a similar number of leaves, more or less similar-sized leaves, and similar plant quality, except that Strawberry Star™ caladium plants are 2–3 inches taller. Tuber de-eyeing seems to be optional for container plant production, but this practice can improve the quality of containerized Strawberry Star™ caladium plants. Strawberry Star™ caladium plants show excellent sunburn tolerance and produce a large number of leaves in sunny locations, thus this variety is suitable for a wide range of locations (shady to sunny) in the landscape.

Availability

Strawberry Star™ caladium was released under the cultivar name 'UF-85-5'. The United States Patent and Trademark Office has issued a Plant Patent (PP24,681) for this cultivar. Production of this variety is to be with a licensing agreement with the Florida Foundation Seed Producers, Inc., PO Box 309, Greenwood, FL 32443. Information on tuber availability and propagation agreements can be obtained from the Florida Foundation Seed Producers, Inc. (<http://ffsp.net/>).

Literature Cited

- Bell, M. L., G. J. Wilfret, and D. A. DeVoll. 1998. "Survey of Caladium Tuber Producers for Acreage of Varieties Grown." *Proceedings of Florida State Horticultural Society* 111: 32–34.
- Deng, Z., and B. K. Harbaugh. 2011. "'UF 85-5': A Spotted Caladium Cultivar for Use in Containers and Sunny Landscapes." *HortScience* 46(9): 1326–1329.
<https://doi.org/10.21273/HORTSCI.46.9.1326>
- Deng, Z., B. K. Harbaugh, R. K. Schoellhorn, and R. C. Andrew. 2008. *2003 Survey of the Florida Caladium Tuber Production Industry*. ENH1007. Gainesville: University of Florida Institute of Food and Agricultural Sciences.
<https://doi.org/10.32473/edis-ep258-2008>
- Wilfret, G. J., and G. T. Hurner, Jr. 1982. "A Survey of Caladium Varieties Grown in Florida and Their Characteristics as Potted Plants." *Proceedings of Florida State Horticultural Society* 95: 190–194.

Tables

Table 1. Tuber weight, production index, marketable number, and grade distribution of Strawberry Star™ caladium and three commercial caladium varieties in 2005 and 2006

Varieties	Tubers			Tuber grade distribution (%)				
	Weight (lb)	Production index	Marketable (no.)	Super Mammoth	Mammoth	Jumbo	No. 1	No. 2
Year: 2005								
Strawberry Star™	12.3	171	50.6	6.3	13.3	30.0	39.3	11.3
Candidum	5.7	103	33.7		4.0	47.7	42.3	6.0
Marie Moir	11.0	159	46.2	2.3	16.7	37.3	35.0	8.7
Miss Muffet	7.1	115	37.3		10.3	40.0	40.0	10.3
Year: 2006								
Strawberry Star™	12.1	175	49.8	7.3	16.0	30.7	29.7	16.3
Candidum	6.6	114	34.6	2.3	11.7	38.7	37.7	9.3
Marie Moir	3.3	58	29.9	2.3	0.0	11.3	43.7	43.0
Miss Muffet	6.0	106	38.7		10.0	28.3	42.7	19.0
<p><i>Note:</i> Values presented are means of three replications with 30 propagules planted in a plot in the field. The production index is an indicator of the economic value of the tubers harvested and is calculated as follows: n (No. 2) + $2n$ (No. 1) + $4n$ (Jumbo) + $6n$ (Mammoth) + $8n$ (Super Mammoth), where n is the number of tubers in each grade.</p> <p>Tubers graded by maximum diameter: No. 2 (1–1.5 inches), No. 1 (1.5–2.5 inches), Jumbo (2.5–3.5 inches), Mammoth (3.5–4.5 inches), and Super Mammoth (> 4.5 inches).</p>								

Table 2. Plant performance and quality of Strawberry Star™ caladium and two commercial varieties grown from No. 1 tubers in 4.5-inch containers in a 45% shaded glasshouse in Wimauma, Florida, in 2007

Variety	Days to sprout		Plant height (inch)		Leaves (no.)		Leaf length (inch)		Leaf width (inch)		Quality rating	
	Intact	De-eyed	Intact	De-eyed	Intact	De-eyed	Intact	De-eyed	Intact	De-eyed	Intact	De-eyed
Strawberry Star™	29.9	32.4	8.8	7.5	12.4	21.2	8.1	6.5	5.5	4.0	3.4	4.4
Marie Moir	34.1	41.2	9.2	7.7	4.9	6.8	11.5	7.3	8.0	4.8	1.9	2.4
Miss Muffet	25.6	30.4	5.8	5.7	10.4	17.7	7.5	5.8	4.8	3.2	3.3	4.1
<p><i>Note:</i> One tuber was planted per container. Data were taken 8 weeks after planting. Values represent the average of 10 plants. "Days to sprout" is the number of days from planting to the first unfurled leaf. Plant quality was rated on a scale of 1–5, with 1 being very poor, 3 being fair and acceptable, and 5 being excellent in plant vigor, fullness, and color display as pot plants.</p>												

Table 3. Plant characteristics, performance, and sunburn tolerance of Strawberry Star™ caladium three commercial varieties, and Cranberry Star™ caladium in 2005 and 2006^Z

Variety	Plant height (inch)	Leaf ^Y			Overall plant performance ratings ^X			Sun tolerance rating ^W		
		number	length (inch)	width (inch)	June	July	August	June	July	August
Strawberry Star™	16.5	27.3	9.8	5.9	3.6	4.3	4.8	3.5	4.5	4.8
Candidum	12.4	14.6	9.4	6.1	1.7	1.5	3.1	2.7	2.6	3.6
Cranberry Star™	14.6	24.9	9.8	6.2	2.6	4.0	4.6	1.8	3.6	3.0
Marie Moir	11.1	13.9	9.6	6.1	1.4	1.6	3.3	1.9	4.2	4.1
Miss Muffet	7.1	16.3	6.6	4.3	1.5	1.3	2.4	3.3	4.4	4.2

^Z Plants were grown from 1-inch tuber propagules in ground beds in full sun in Wimauma, Florida. Values for plant height, leaf number, length, and width are means of three replications with three plants measured per plot per year, while performance and sunburn tolerance ratings are means of three replications based on whole plot evaluation.

^Y Data were taken over two growing seasons (2005 and 2006), approximately 4 months (August 2005 and 2006) after tubers were planted in April each year.

^X Plants were rated on a scale of 1–5, with 1 being very poor, 3 being fair and acceptable, and 5 being excellent in plant vigor, fullness, and color display, in June, July, and August 2005 and 2006.

^W Plant sunburn tolerance was rated on a scale of 1–5, with 1 being very poor, 3 being fair and acceptable, and 5 being excellent without showing any signs of sunburn or resulting holes on leaf surfaces, in June, July, and August 2005 and 2006, respectively.

¹ This document is ENH1201, one of a series of the Environmental Horticulture Department, UF/IFAS Extension. Original publication date August 2012. Revised September 2015. Visit the Ask IFAS website at <https://ask.ifas.ufl.edu> for the currently supported version of this publication.

² Zhanao Deng, professor, breeding and floriculture, Department of Horticultural Sciences, UF/IFAS Gulf Coast Research and Education Center; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information, and other services only to individuals and institutions that operate in compliance with applicable federal and state non-discrimination laws and policies. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A&M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, Dean for UF/IFAS Extension.