

Malus x 'Jewelberry' Jewelberry Crabapple¹

Edward F. Gilman, Ryan W. Klein, and Gail Hansen²

Introduction

Crabapples are best grown in a sunny location with good air circulation and have no particular soil preferences, except soil should be well-drained. Root-pruned trees transplant most easily. Tree size, flower color, fruit color, and growth and branching habit vary considerably with the cultivar grown, but many grow about 20 feet tall and wide. A few crabapples have good fall color, and double-flowered types hold blossoms longer than single-flowered cultivars. Some crabapples are alternate bearers, blooming heavily only every other year. Crabapples are grown for their showy flowers and attractive, brightly colored fruit.



Figure 1. Full Form - *Malus x 'Jewelberry'*: Jewelberry Crabapple
Credits: Edward F. Gilman, UF/IFAS



Figure 2. Leaf - *Malus x 'Jewelberry'*: Jewelberry Crabapple
Credits: Edward F. Gilman, UF/IFAS



Figure 3. Fruit - *Malus x 'Jewelberry'*: Jewelberry Crabapple
Credits: Edward F. Gilman, UF/IFAS

1. This document is FPS-392, one of a series of the Department of Environmental Horticulture, UF/IFAS Extension. Original publication date October 1999. Revised October 2023. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. Edward F. Gilman, professor emeritus; Ryan W. Klein, assistant professor, arboriculture; and Gail Hansen, professor, sustainable landscape design; Department of Environmental Horticulture, UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, dean for UF/IFAS Extension.

General Information

Scientific name: *Malus x 'Jewelberry'*

Pronunciation: MAY-luss

Common name(s): 'Jewelberry' crabapple

Family: *Rosaceae*

Plant type: tree

USDA hardiness zones: 4 through 8A (Figure4)

Planting month for zone 7: year round

Planting month for zone 8: year round

Origin: not native to North America

Invasive potential: not known to be invasive

Uses: specimen; espalier; trained as a standard; recommended for buffer strips around parking lots or for median strip plantings in the highway; bonsai

Availability: somewhat available, may have to go out of the region to find the plant



Figure 4. Shaded area represents potential planting range.

Description

Height: 8 to 12 feet

Spread: 10 to 15 feet

Plant habit: round

Plant density: dense

Growth rate: moderate

Texture: medium

Foliage

Leaf arrangement: alternate

Leaf type: simple

Leaf margin: serrate

Leaf shape: elliptic (oval)

Leaf venation: brachidodrome

Leaf type and persistence: deciduous

Leaf blade length: 2 to 4 inches

Leaf color: green

Fall color: yellow

Fall characteristic: not showy

Flower

Flower color: pink

Flower characteristic: showy

Fruit

Fruit shape: round

Fruit length: less than 0.5 inch

Fruit cover: fleshy

Fruit color: red

Fruit characteristic: attracts birds; showy; persists on the plant

Trunk and Branches

Trunk/bark/branches: typically multi-trunked or clumping stems; not particularly showy; no thorns

Current year stem/twig color: brown

Current year stem/twig thickness: medium

Culture

Light requirement: plant grows in full sun

Soil tolerances: clay; acidic; well-drained; sand; loam; alkaline; occasionally wet

Drought tolerance: unknown

Soil salt tolerances: moderate

Plant spacing: not applicable

Other

Roots: usually not a problem

Winter interest: plant has winter interest due to unusual form, nice persistent fruits, showy winter trunk, or winter flowers

Outstanding plant: not particularly outstanding

Pest resistance: long-term health usually not affected by pests

Use and Management

Plants are used for specimens, patios (small fruiting types), and along streets to create a warm glow of color each spring. Most are attractive during the summer, bearing glossy green foliage. Popular around overhead power lines due to their small stature, a row of crabapples along each side of the street or median strip can “make” a neighborhood. Select plants which have been grafted onto EMLA 106 or 111 rootstock to reduce root suckering.

Crabapple is well-adapted to compacted urban soil, tolerates drought and poor drainage well, and is somewhat tolerant of salt-spray. It is well-adapted to all areas within its hardiness zone range, including Texas and Oklahoma. Very adaptable tree. Do not overfertilize, since this could increase the incidence of disease. Select only from disease-resistant cultivars. Crabapples grow well in the Texas panhandle but are not extremely drought tolerant and are not well suited for high-pH soil.

Large-fruited types can create a maintenance problem because rotting fruits attract insects and rodents and are quite messy. Some crabapples sprout vigorously from the roots, and these will require regular pruning to maintain an attractive tree. Trees used as street trees will require regular pruning early in their lives to train lower branches for pedestrian and vehicle clearance.

There are hundreds of crabapple cultivars with single or double red, pink, or white flowers, and varying fruit size. Many are cultivars of *Malus baccata* and *Malus floribunda*.

Since disease resistance can vary depending on where a particular cultivar is grown, be sure to choose a cultivar that has been shown to be resistant to disease in your area. Your urban forestry program could suffer if you plant the wrong cultivar, but it could blossom if the correct ones are installed.

Malus adapted for street tree and urban use include ‘Adams’, ‘Bob White’, ‘David’, ‘Donald Wyman’, ‘Profusion’, ‘Red Splendor’ and *Malus floribunda*. Be sure to specify tree form plants for street tree use since branching may be too low on trees grown for specimen use.

White - flowered cultivars include ‘BaccataColumnaris’—narrow crown, white flowers, red or yellow fruit; ‘Baccata Gracilis’—slow-growing, shrub-like, white flowers, fruit small and dark red, annual bearer; ‘BaccataJackii’—upright form, white flowers, bright red fruit, annual bearer, also good to excellent disease resistance; ‘Callaway’—pink buds, white flowers, red fruit; ‘David’—pink buds open to white flowers, scarlet fruit, good to excellent disease resistance; ‘Dolgo’—pink buds, white flowers, large red fruits; ‘Donald Wyman’—disease-resistant but susceptible to fire blight, glossy red showy fruit; ‘Ellwangeriana’—red fruit, disease-resistant; ‘Floribunda’—pink to red bud opens to single white flower, yellow, or red fruit, commonly available; ‘Gloriosa’—pink bud opens to white flower, red, large fruit; ‘Golden Hornet’—upright arching habit, white flower, yellow fruit; ‘Gorgeous’—pink bud opens to large, white flower, red to orange fruit; ‘Harvest Gold’—white flowers followed by yellow fruits; ‘Hupehensis’—tea crabapple—pink buds open to white flowers, greenish fruit; ‘Katherine’—double flowers opening pink, fading to white, fruit yellow and red; ‘Mary Potter’—pink buds open to single white flowers, red and fairly large fruit, susceptible to scab and powdery mildew; ‘Red Jade’—weeping habit, white flowers, red fruit persisting after leaves drop; ‘Sargentii’—dwarf, pink bud opens to white flowers, small dark red fruit; ‘Snowdrift’—white flowers, orange - red fruit; ‘Tanner’—white flowers, red fruits, susceptible to diseases; ‘Tschonoski’—white flowers, vigorous growth, good bronze red fall color, fruit brownish; ‘White Angel’—white flowers, glossy red fruit persisting into winter; ‘White Candle’—pink buds open to white flowers, red fruit, upright growth habit; ‘ZumiCalocarpa’—white flowers, bright red, persistent fruit.

Pink-flowered cultivars include ‘Arnoldiana’—red buds, pale pink to white flowers, red and yellow fruits, susceptible

to diseases; 'Barbara Ann'—double pink flowers; 'Beverly'—pink flowers, small, bright red fruit, upright open habit; 'Brandywine'—double, pink flowers, foliage tinged reddish; 'Candied Apple'—red buds and pink flowers, persistent, bright red fruits; 'Coralburst'—pink flowers, a dwarf about eight feet tall; 'Dorothea'—red buds, pink flowers, yellow fruits, susceptible to scab; 'Hopa'—pink flowers, red and yellow fruit, extremely susceptible to scab; 'Margaret'—disease-resistant; 'Mount Arbor Special'—disease-resistant, red fruit; 'OekonomierathEchtermayer'—somewhat weeping habit, red buds, pink flowers, red fruits; 'Pink Perfection'—pink flowers, susceptible to scab; 'Prairie Rose'—double pink flowers, no fruit; 'Radiant'—red buds, pink flowers, persistent red fruits, susceptible to scab; 'Van Eseltine'—double pink flowers, columnar; 'Velvet Pillar'—upright form, pink flowers, red fruit and purple foliage; 'Weeping Candied Apple'—pink flowers and red fruit on horizontal to pendulous branches.

Cultivars with red flowers include 'American Beauty'—double, red flowers, susceptible to scab; 'Centurion'—a columnar growth habit, red flowers followed by bright red, persistent fruits; 'Indian Magic'—red flowers, maroon fall color, small red fruit, persistent; 'Purpurea Lemoinei'—dark rose color flowers, alternate bearing; 'Profusion'—red flowers and fruit; 'Red Baron'—red flowers, susceptible to scab; 'Red Silver'—red flowers, red fruit, susceptible to scab; 'Red Splendor'—red buds, red to pink flowers, red fruit, susceptible to scab; 'Royal Ruby'—double red flowers, sparse red fruit, susceptible to scab; 'Royalty'—red flowers, red fruits, young foliage purple, susceptible to scab; 'Selkirk'—red flowers, red fruits, susceptible to diseases; 'Sparkler'—red flowers, red fruits; 'Tomiko'—disease-resistant, reddish-purple blooms; 'Wisley'—red flowers.

One of the best crabapples for the south is *Malus x Callaway*.

Disease-resistant cultivars include: 'David', 'Dolga', 'Donald Wyman', 'Ellwangeriana', 'Inglis', 'Jackii', 'Jewelberry', 'Margaret', 'Mary Potter', 'Mount Arbor Special', 'Prairifire', 'Professor Sprenger', 'Tomiko'. Contact the Ornamental Crabapple Society, Morton Arboretum, Lisle, Illinois 60532, for more information on crabapples.

Pests and Diseases

Aphids infest branch tips and suck plant juices and are quite common. They can deform newly emerging foliage, and they secrete honeydew, creating a sticky mess beneath the tree, but they will not kill the tree.

Fall webworm makes nests on the branches and feeds on foliage inside the nest. Small nests can be pruned out or sprayed with *Bacillus thuringiensis*. Controlling severe infestations may require other chemicals.

Scales of various types are controlled with horticultural oil.

Borers can be a problem on stressed trees.

Mites are too small to be seen easily, so they can cause much foliage discoloration before being detected. Mites can be controlled to a degree with horticultural oil, but other chemicals are often required by the time mites are detected. The mite infestation can also be severe by the time foliage chlorosis or bronzing is evident.

Eastern tent caterpillar builds tents or nests in trees in early summer or late spring. Feeding occurs on foliage outside the nest. Defoliation can be extensive if infestation is severe, and repeated defoliations for several years can weaken trees. Small nests can be removed by pruning them from the tree. Spray with *Bacillus thuringiensis* or another approved chemical. Do not burn nests while they are still in the tree.

Many selections are fairly susceptible to disease. Select from the disease-resistant ones.

Scab infection takes place early in the season and dark olive-green spots appear on the leaves. In late summer the infected leaves fall off when they turn yellow. Infected fruits have black, slightly raised spots. Use resistant varieties to help avoid this severe problem.