

# The Horntail Snail (*Macrochlamys indica*): A New Invasive Pest in Florida<sup>1</sup>

Alexandra M. Revynthi, Daniel Carrillo, Dakshina R. Seal, E. Vanessa Vassilaros, and Paul E. Kendra<sup>2</sup>

## Introduction

The purpose of this document is to provide an overview of the current knowledge on the horntail snail (*Macrochlamys indica* Benson) as well as recommendations to optimize its detection and monitoring in nurseries.

The horntail snail was detected in August 2020 in Miami-Dade County (Talamas 2020). This snail is considered of quarantine importance in the United States because it can become an agricultural pest and potentially host parasitic nematodes of medical significance (Grewal et al. 2003; Cowie et al. 2009; Jayashankar and Murthy 2015). A program to survey, control, and eradicate the horntail snail has been implemented by the Florida Department of Agriculture and Consumer Services, Division of Plant Industry (FDACS-DPI).



Figure 1. Worldwide distribution of the horntail snail (*Macrochlamys indica* Benson).

Credit: undefined

The horntail snail belongs to the genus *Macrochlamys* (family Ariophantidae), which has over a hundred described species distributed from South to Southeast Asia and southern China (Pholyotha et al. 2018). It occurs in India, Sri Lanka, Bangladesh, Nepal, Pakistan and Brazil (Figure 1) (Raut and Ghose 1984; Biswas et al. 2015; Jayashankar et al. 2015; Agudo-Padron 2018). Jayashankar et al. (2015) report the occurrence of this snail in Europe; however, the location is not specified, and we were unable to find additional information to substantiate this report. In Bangalore, India, it is known as the common garden snail, and populations are higher

in urban settings than in agricultural areas (Jayashankar et al. 2015).

## Identification

The horntail snail has a brown body and an amber-colored shell (Figure 2). At the tip of its tail, *M. indica* has a caudal horn, from which it took its common name (Talamas 2020). The size of adult snails varies with age, and the shell diameter can range from 10 to 21 mm (0.39 – 0.83 in) (Raut and Ghose 1984) (Table 1). A characteristic of this snail species is that it has a flap of flesh that extends backward onto or around the shell (for pictures, see Talamas 2020). In Florida, the jump snail (*Ovachlamys fulgens* (Gude)) has similar characteristics, such as a caudal horn and similar body and shell coloration. However, this snail has a thinner and more elongated body, and its caudal horn is much larger and thicker (for description and pictures, see Salles et al. 2018).



Figure 2. The horntail snail (*Macrochlamys indica* Benson).

Credit: FDACS-DPI

## Host range and damage

Information is scarce regarding the pest and vector status of the horntail snail in its native range. It feeds mostly on seedlings and fallen, decomposed leaves (Raut and Ghose 1983, 1984; Jayashankar et al. 2015). It is considered a minor pest in some localities in India, where it feeds almost exclusively on moss growing on damp brick walls (Jayashankar et al. 2015). The horntail snail, however, can feed on various plants, including vegetable, fruit, and ornamental crops. In caged studies,

preferred vegetables included cucurbits, beans, lettuce, okra, and cabbage, while papaya was the preferred fruit crop. Ornamental plants such as chrysanthemum, china rose hibiscus, marigold, and roses were also considered preferred hosts in caged studies (Raut and Ghose 1983, 1984). In general, younger snails feed on tender plant tissue and avoid the leaf petioles and veins, while adults consume all plant parts (Raut and Ghose 1983).

## Biology

The biology of horntail snails was studied under laboratory and field conditions in Bangalore, India. Relative humidity (RH) was found to be the most important factor affecting the population dynamics of this snail species (Raut and Ghose 1979, 1984). When the humidity in the environment is below 46%, the horntail snail enters a dormant state called aestivation. Aestivation varies with geographic latitude and can range from two to eight months. During this period, the snails remain hidden and inactive and lose weight (Raut and Ghose 1979, 1984). In southern India, the horntail snail has an eight-month aestivation period and remains active during the monsoon season. A single individual can live up to four years. Young snails require 130 days to reach sexual maturity. Optimum mating conditions consist of temperatures ranging from 21°C–30°C (69.8°F–86°F) and humidity higher than 86%. Mating usually takes place after rains, and the gestation period ranges from 10–17 days.



2nd year.



3rd year.



1st year.



4th year.

In India, egg laying starts when snails are in their second year (Table 1). Eggs are laid in clutches under loose soil. Adult snails need approximately 10–24 minutes to deposit an egg clutch and, on average, remain in the nest 11 minutes after that (Raut and Ghose 1984). Eggs are round, translucent, and 3.0–3.5 mm (0.12–0.14 in) in size. The number of egg clutches and eggs per clutch varies with the age of the female (see Table 1). The egg clutches are laid at 12 to 49-day intervals, and the egg incubation period ranges from eight to twelve days. After egg hatching, young snails remain immobile for 14–20

hours, and they start feeding on the fourth or fifth day (Raut and Ghose 1984). The snail, however, might not have the same reproductive and feeding habits in new areas of invasion.

Under laboratory conditions with temperature ranging from 21°C to 32°C (69.8°F–89.6°F) and constant 90% RH, the horntail snail's life span ranges from 15 to 19 months, and there is no aestivation. Snails oviposit up to five egg clutches per year with an average of 16 eggs per clutch. Under these conditions, young snails require ten months to reach sexual maturity (Sudhakar 2013).

## Dispersal and Environment

Like other land snails, the horntail snail has low natural long-distance dispersal. Human transport of plants and landscape materials provides the major mechanism for introduction of snails into new locations (Bergey et al. 2014). The low long-distance mobility of land snails also offers the opportunity to eradicate local populations (Roda et al. 2016). As in most land snails, active dispersal occurs at night and on cloudy, rainy days (Bailey 1981).

Relative humidity is the most critical factor affecting the biology and behavior of the horntail snail. Temperature ranging from 28°C to 31°C (82.4°F –87.8°F) with a minimum of 19°C (66.2°F) and relative humidity between 85 and 100% are the optimum environmental conditions for the development of this species. Even when the temperature ranges from 9°C to 43°C (48.2°F –109.4°F), the horntail snail can remain active as long as the relative humidity is above 60% (Raut and Ghose 1984).

## Association with Human Pathogens

The association between horntail snail and nematodes has not been studied in detail. A single study mentioned an association with non-parasitic *Rhabditis* sp. nematodes in West Bengal, India (Cowie et al. 2009; Jayashankar and Murthy 2015). However, closely related species, such as *Macrochlamys resplendens* and the Giant African Snail (*Achatina fulica*), are known hosts of *Angiostrongylus* nematodes that can cause eosinophilic meningitis in humans (Grewal et al. 2003). Therefore, the possibility that the horntail snail could host nematodes of medical importance cannot be ruled out.

## Recommendations to Optimize Detection and Monitoring of Horntail Snail Populations

Early detection and monitoring of horntail snails in the field are crucial for its successful control/eradication. Recommendations to optimize detection of snail populations include:

1. Search for snails at night when they are active, especially after rainfall. Remember that there are

also other snails present in our environment. Proper identification is key for a successful management program. FDACS-DPI can assist with snail identification.

2. During the day, search for snail trails, shells, and plant damage.
3. Search for snails in debris piles, bricks, wall crevices, and near seedlings or succulent plants.
4. Search for snails in areas that remain moist and humid, such as near air-conditioning units, pump irrigation systems, and under mulch.
5. Train personnel working on ornamental, vegetable, and tropical fruit production on when and how to monitor for snail populations.

In addition, the following actions can be taken to reduce snail populations:

1. Improve sanitation and remove hiding places such as boards, stones, debris, weedy areas, leafy branches growing close to the ground, dense ground covers and mulch, and any areas that remain moist.
2. If possible, create a less humid environment by adjusting the irrigation frequency and intensity, promptly repairing leaks, and improving drainage.

If you find the horntail snail on your property contact your local county UF/IFAS Extension office (see <http://sfyl.ifas.ufl.edu/find-your-local-office/>) and the Florida Department of Agriculture and Consumer Services – Division of Plant Industry (FDACS-DPI). For more information from DPI helpline, go to <https://www.fdacs.gov/Agriculture-Industry/Pests-and-Diseases/Plant-Pests-and-Diseases/Invasive-Mollusks>, call 1-888-397-1517 or email photos to [DPIhelpline@FDACS.gov](mailto:DPIhelpline@FDACS.gov). Properties where the horntail snail has been detected are required to hold orders and quarantine for at least 30 days, sign a compliance agreement, and follow the treatment protocol with metaldehyde (>3.25%) according to FDACS recommendations. For more information, go to the current FDACS Pest Alert document: <https://www.fdacs.gov/content/download/93400/file/horntail-snail-pest-alert.pdf>. Registered molluscicides for Florida with metaldehyde (>3.25%) for use in nurseries and ornamental crops can be found in Table 2. Note: Do not touch snails with bare hands. Gloves are advised when handling the snail.

## References

- Agudo-Padron, A. I. 2018. "Revised And Updated Systematic Inventory of Non-Marine Molluscs Occurring in the State of Santa Catarina/S.C., Central Southern Brazil Region." *Advances in Environmental Studies* 2:74–81
- Bailey, S. E. R. 1981. "Circannual and Circadian Rhythms in the Snail *Helix aspersa* Müller and

- the Photoperiodic Control of Annual Activity and Reproduction." *Journal of Comparative Physiology* 142:89–94.  
<https://doi.org/10.1007/BF00605480>
- Bergey, E. A., L. L. Figueroa, C. M. Mather, R. J. Martin, E. J. Ray, J. T. Kurien, D. R. Westrop, and P. Suriyawong. 2014. "Trading in Snails: Plant Nurseries as Transport Hubs for Non-Native Species." *Biological Invasions* 16:1441–1451.  
<https://doi.org/10.1007/s10530-013-0581-1>
- Biswas, T., B. Tripathy, K. Valarmathi, and S. K. Sajan. 2015. "Taxonomy, Distribution and Conservation of Molluscs in Kangra District of Himachal Pradesh: Three New Records from the State." *Ambient Science* 2:18–24.  
<https://doi.org/10.21276/ambi.2015.02.2.ra02>
- Cowie, R. H., R. T. Dillon, D. G. Robinson, and J. W. Smith. 2009. "Alien Non-Marine Snails and Slugs of Priority Quarantine Importance in the United States: A Preliminary Risk Assessment." *American Malacological Bulletin* 27:113–132.  
<https://doi.org/10.4003/006.027.0210>
- Grewal, P. S., S. K. Grewal, L. Tan, and B. J. Adams. 2003. "Parasitism of Molluscs by Nematodes: Types of Associations and Evolutionary Trends." *Journal of Nematology* 35:146–156
- Jayashankar, M., and G. S. S. Murthy. 2015. "Record of Gut Associated Nematelminth in the Giant African Snail *Achatina fulica* (Bowdich) from Bangalore, India." *Journal of Parasitic Diseases* 39:144–146. <https://doi.org/10.1007/s12639-013-0303-8>
- Jayashankar, M., M. S. Reddy, and S. Ramakrishna. 2015. "Incidence of the Common Garden Snail, *Macrochlamys indica* Benson, 1832 (Gastropoda: Ariophantidae) in Bangalore Region." *The Bioscan* 10:1003–1006
- Pholyotha, A., C. Sutcharit, and S. Panha. 2018. "The Land Snail Genus *Macrochlamys* Gray, 1847 from Thailand, with Descriptions of Five New Species (Pulmonata: Ariophantidae)." *Raffles Bulletin of Zoology* 7600:763–781
- Raut, S. K., K. C. Ghose. 1984. "Pestiferous land snails of India." *Zoological Survey of India* 1–151
- Raut, S. K., and K. C. Ghose. 1983. "Food Preference and Feeding Behaviour of Two Pestiferous Snails, *Achatina fulica* Bowdich and *Macrochlamys indica* Godwin-Austen." *Records of the Zoological Survey of India* 80:421–440
- Raut, S. K., and K. C. Ghose. 1979. "Factors Influencing Mortality in Land Snails, *Achatina fulica* Bowdich and *Macrochlamys indica* Godwin-Austen during Aestivation." *Proceedings of the Zoological Society of London* 32:107–120
- Roda A, G. Nachman, S. Weihman, M. Y. Cong, and F. Zimmerman. 2016. "Reproductive Ecology of the Giant African Snail in South Florida: Implications for Eradication Programs." *PLoS One* 11:1–18.  
<https://doi.org/10.1371/journal.pone.0165408>
- Salles, A. C. A., C. D. C. Oliveira, and R. S. Absalão. 2018. "Redescription of the Jumping Snail *Ovachlamys fulgens* (Glide, 1900) (Gastropoda: Helicarionoidea: Helicarionidae): An Anatomical and Conchological Approach." *Nautilus* (Philadelphia) 132:19–29
- Sudhakar, B. S. 2013. "Chapter 3, Life Cycle of *Macrochlamys indica*." *Effect of thiamethoxam and diafenthiuron on land snail Macrochlamys indica (Bensonpulmonata: Ariophantidae)*. Ph.D. Dissertation, North Maharashtra University: 42–75
- Talamas, E. J. 2020. Pest Alert: Horntail Snail, *Macrochlamys indica* Benson.

## Tables

Table 1. In India, under field conditions, horntail snails (*Macrochlamys indica* Benson) of different ages vary in size and oviposition rates (Raut and Ghose 1984). Photo credit: FDACS-DPI.

Snail age	Shell diameter (mm)	Number of egg clutches/ breeding season	Av. Number of eggs per clutch
1st year	10–11 (0.39–0.43 in)	NA	NA
2nd year	10–14 (0.39–0.55 in)	4	36
3rd year	14–18 (0.55–0.71 in)	6	69
4th year	18–21 (0.71–0.83 in)	4	89

Table 2. Molluscicides registered in Florida for use in nurseries and ornamental crops. Products contain more than 3.25% metaldehyde, the minimum percentage recommended by FDACS for eradication of the horntail snail (*Macrochlamys indica* Benson).

Active ingredient	Brand name	Rate	Max. number of applications/growing season	Restricted-Entry Interval REI (hrs)	Reapplication interval (days)
Metaldehyde 3.5%	Durham Metaldehyde 3.5%	1 lb a.i. / Acre  Max 2 lb a.i. for eradication	6	12	21
Metaldehyde 4%	Deadline bullets	25 lbs / Acre  Max 50 lbs / Acre for eradication	6	12	21
	Deadline M-Ps	25 lbs / Acre  Max 50 lbs / Acre for eradication	6	12	21
	Durham ornamental	24 pellets/ sq ft	6	NA	21
	Metarex	8–25 lbs / Acre (6–20 pellets / sq ft)	6	12	21
	Sluggo 4.0	25 lbs / Acre	6	NA	21
Metaldehyde 7.5%	Durham Metaldehyde 7.5%	1 lb a.i. / Acre	6	12	21
Metaldehyde 25%	Slug Fest All weather formula	59 fl oz / Acre	6	12	21

<sup>1</sup> This document is ENY2078, one of a series of the Department of Entomology and Nematology, UF/IFAS Extension. Original publication date April 2022. Please visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.

<sup>2</sup> Alexandra M. Revynthi, assistant professor, agricultural acarology, Department of Entomology and Nematology, UF/IFAS Tropical Research and Education Center, Homestead, FL; Daniel Carrillo, associate professor, Department of Entomology and Nematology, UF/IFAS Tropical Research and Education Center; Dakshina R. Seal, scientist, pest management, UF/IFAS Tropical Research and Education Center, Homestead, FL; E. Vanessa Vassilaros, former Extension agent III, M.S., commercial agriculture/ornamentals, UF/IFAS Extension Miami-Dade County, Homestead, FL; Paul E. Kendra, research entomologist, Subtropical Horticulture Research, USDA Agricultural Research Service; UF/IFAS Extension, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition.

Use pesticides safely. Read and follow directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, dean for UF/IFAS Extension.