

Asian Woolly Hackberry Aphid, *Shivaphis celti* Das (Insecta: Hemiptera: Aphididae)¹

Susan E. Halbert and Paul M. Choate²

The Featured Creatures collection provides in-depth profiles of insects, nematodes, arachnids and other organisms relevant to Florida. These profiles are intended for the use of interested laypersons with some knowledge of biology as well as academic audiences.

Introduction

An Asian woolly hackberry aphid, *Shivaphis celti* Das, was found for the first time in Florida in Jacksonville, Duval County, on sugarberry (*Celtis laevigata* Willd.) on 13 August 1997 by Florida Department of Agriculture & Consumer Services' Division of Plant Industry Plant Inspector Flewellyn W. Podris. Since then, these Asian woolly hackberry aphids have been collected in counties spanning most of Florida. *Shivaphis celti* was found in Georgia about a year before its discovery in Florida.

There are several species of Asian woolly hackberry aphids. The East and Central Asian genus *Shivaphis* contains six described species, of which four are in *Shivaphis sensu stricto*, and the other two are in the subgenus *Shivaphis (Sinishivaphis)* (Quednau and Remaudière 1985; Remaudière and Remaudière 1997; Zhang and Zhong 1982). They are found primarily on *Celtis* spp., with one described species, *Shivaphis (Sinishivaphis) tilisucta* Zhang, from China on *Tilia* (Zhang and Zhong 1990). Two more species, included in *Shivaphis* in older literature, are now in the

genus *Neocranaphis* (Remaudière and Remaudière 1997). These species infest bamboo and closely related plants. No Asian woolly hackberry aphids other than *S. celti* are known to occur in the Western Hemisphere.

Description

Shivaphis celti is conspicuous because the aphids secrete copious quantities of bluish white wax. The aphids are small, about 2 to 2.5 mm long. As is often the case with Myzocalidini, the anal plate is deeply cleft. The cauda is finger-like rather than constricted into a knob. There are conspicuous wax glands on the abdomen. Siphunculi are pore-like, on a slightly raised cone. Wings veins are bordered, particularly at the distal ends. Antennal joints are darkened, giving the antenna a striped appearance. The processus terminalis is rather short, about three times as long as wide. These characters will separate *S. celti* from other genera of aphids on *Celtis*, and from other Asian woolly hackberry aphids (Blackman and Eastop 1994; Quednau and Remaudière 1985).

1. This document is EENY288 (originally published as DPI Entomology Circular 392), one of a series of the Entomology and Nematology Department, UF/IFAS Extension. Original publication date June 2003. Revised March 2021. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication. This document is also available on the Featured Creatures website at <http://entnemdept.ifas.ufl.edu/creatures/>.
2. Susan E. Halbert, Division of Plant Industry, Florida Department of Agriculture and Consumer Services; and Paul M. Choate, Entomology and Nematology Department; UF/IFAS Extension, Gainesville, FL 32611.



Figure 1. Winged adult female *Shivaphis celti* Das, an Asian hackberry aphid, on hackberry.

Credits: P. M. Choate, UF/IFAS

Life History

Summer adults are all female and parthenogenetic. Summer adults may be winged or wingless. In the autumn (October in Gainesville), winged males and wingless oviparae can be found. These mate to produce an overwintering egg that allows the aphids to survive the winter when there are no leaves on the trees. Raychaudhuri et al. (1981) described the oviparous forms.



Figure 2. Wingless adult *Shivaphis celti* Das, an Asian hackberry aphid, on hackberry.

Credits: P. M. Choate, UF/IFAS

Hosts

Shivaphis celti probably is restricted to *Celtis*, although Chakrabarti (1988) found *S. celti* occasionally on *Arundo donax* L. Host species of *Celtis* include: *C. australis* L., lote-tree, Mediterranean hackberry (Chakrabarti 1988; Raychaudhuri et al. 1980); *C. jessonensis* Koidz, Korean hackberry/ezo-enoki (Quednau, 1979); *C. laevigata* (Florida observations); *C. nervosa* Hemsl., little leaf hackberry (Chinese common name) (Zhang and Zhong 1983); *C. sinensis* Pers., Japanese hackberry (Higuchi 1972; Zhang

and Zhong 1983); *C. tetrandra* Roxb., Malayan hackberry, (Chakrabarti 1988); and *C. tetraneura* [sic] (Raychaudhuri et al. 1980). Other hackberry species listed only by Chinese common name include: green hackberry, sand or desert hackberry, Yunnan hackberry, American hackberry and Australian hackberry (Zhang and Zhong 1983).

Survey and Detection

Look for small (up to 2 mm) balls of fuzzy wax on the backs of hackberry leaves. Infestations can be found in both urban and natural landscape situations.



Figure 3. A colony of *Shivaphis celti* Das, an Asian hackberry aphid, colony on *Celtis*. Note copious quantities of bluish white wax around the insects.

Credits: L. J. Buss, UF/IFAS

Management

Chemical

No long-term damage associated with woolly hackberry aphids has been observed; however, aphids may become quite numerous on some trees in late summer. Chemical treatment probably is not warranted for protecting the health of the infested trees.

Biological

Two aphid parasitoids were described from *S. celti*, including *Trioxys (Trioxys) soporensis* Shujauddin (Shujauddin 1982) and *Trioxys (Binodoxys) jaii* Bhagat (Bhagat 1982). Another parasitoid, *Trioxys (Trioxys) shivaphis* Takada, is mentioned in Shujauddin (1982). The efficacy of these parasitoids in controlling the population is not known. No parasitized *S. celti* have been found in Florida so far.

Selected References

- Bhagat RC. 1982. On two new aphid parasitoids of genus *Trioxys* (Aphidiidae: Hymenoptera) from Kashmir, India. *Entomon* 7: 321–324.
- Blackman RL, Eastop VF. 1994. Aphids on the world's trees. CAB International, Wallingford, UK. 466 p.
- Chadrabarti S. 1988. Revision of the Drepanosiphinae (Homoptera: Aphididae) from the Indian subregion. *Oriental Insects* 22: 1–86.
- Higuchi H. 1972. A taxonomic study of the subfamily Callipterinae in Japan. *Insecta Matsumurana* 35: 19–126.
- Quednau FW. 1979. A list of the Drepanosiphine aphids from the Democratic People's Republic of Korea with taxonomic notes and descriptions of new species (Homoptera). *Annales Zoologici Polska Akademia Nauk Instytut Zoologii* 34: 501–528.
- Quednau FW, Remaudière G. 1985. Une nouvelle espèce du genre *Shivaphis* Das de Turquie (Homoptera: Aphididae). *Canadian Entomologist* 117: 227–232.
- Raychaudhuri DN, Ghosh AK, Basu RC, Ghosh MR, Chatterjee M, Chakrabarti S, Pal PK. 1980. Aphids of north-east India and Bhutan. The Zoological Society, Calcutta. 521 p.
- Raychaudhuri D, Raychaudhuri DN, Singh TK. 1981. Redescription of *Shivaphis celti* (oviparae) and *Myzus varians* (Homoptera: Aphididae) hitherto unknown from India. *Science and Culture* 47: 171–172.
- Remaudière G, Remaudière M. 1997. Catalogue des Aphididae du monde. Institut National de la Recherche Agronomique, Paris. 473 p.
- Shujauddin. 1982. Description of a new species of the genus *Trioxys* Haliday (Hymenoptera: Aphidiidae) and new record of *Trioxys* (*Trioxys*) *pallidus* (Hal.) from Kashmir (India). *Journal of Entomological Research* 6: 146–149.
- Zhang G-x, Zhong T-s. 1990. New species and a new record of Callaphididae and Aphididae from northeast China (Homoptera: Aphididae). *Acta Entomologica Sinica* 33: 84–88.
1983. Economic Insect Fauna of China, Homoptera: Aphidinea. Science Press, Beijing. Volume 25, Part I. 387 p.