

Woody Goldenrod, *Chrysoma pauciflosculosa*¹

Debbie Miller, Mack Thetford, Christina Verlinde, and Gabriel Campbell²

Note: This fact sheet is also available as a chapter in a comprehensive manual titled Dune Restoration and Enhancement for the Florida Panhandle, available in pdf form here:

<https://edis.ifas.ufl.edu/pdffiles/SG/SG15600.pdf>. Please see the manual for more information about other useful and attractive native plants for dunes and for further information about restoration and preservation techniques.

Asteraceae



Figure 1. Credit: Mack Thetford, UF/IFAS

Woody goldenrod is found in beach dunes, scrub, and sandhill plant communities. It occurs in the Panhandle of Florida and more broadly in the southeastern United States west to Mississippi and northeast to North Carolina. This plant may leach chemicals into the soil that inhibit seed germination of gulf bluestem (*Schizachyrium scoparium* var. *scoparium*) (Fischer et al. 1994). Woody goldenrod is an underused landscape plant with a prolific fall color from flowers that attract numerous pollinators.

General Description

Woody goldenrod is a multibranched, evergreen subshrub that can reach heights of 3.3 ft or more. Stems are glabrous, numerous, ascending, and round in cross section. Stems bearing flowers and fruits extend far beyond vegetative stems. Leaves are sessile (without stalks), simple, 0.75 to 2.5 in long, alternate, elliptic to oblong, and grayish green with an entire margin. Inflorescences are heads that are sticky, borne in clusters, and numerous, with a single outer row of yellow

ray florets. They occur from late summer to fall. Fruits are hard, dry, pale, bristled achenes.

Propagation

Cutting propagation of woody goldenrod is easily accomplished with terminal stem cuttings. Nonbranched, vegetative apical stem cuttings from summer softwood or winter hardwood root without auxin, but a 5,000 ppm IBA (Indole-3-butyric acid) application will increase root numbers and root length (Wilson et al. 2010). Propagation from seed for woody goldenrod has been accomplished with wild-collected seed. Seeds prefer cooler temperatures (20/10°C alternating day/night temperatures) to warmer ones (Wilson et al. 2010).

Outplanting

Rooted cuttings or seedlings may be grown in a variety of coarse production substrates and a variety of container sizes. Smith et al. (2014) produced woody goldenrod in substrates such as Atlas 3000, Atlas 7000, Fafard 3B, and Metro Mix 300 while the authors have grown them in a 100% pine-bark substrate amended with lime and slow-release fertilizer. Plants grown in 1-gal containers were evaluated for potential use in home landscapes with high transplant success (Smith et al. 2014). Peak flowering for the trial plants occurred in November.

Literature Cited

Fischer, N.H., G.B. Williamson, J.D. Weidenhamer, and D.R. Richardson. 1994. "In search of allelopathy in the Florida scrub: the role of terpenoids." *Journal of Chemical Ecology* 20(6)1355–1380. <https://doi.org/10.1007/BF02059812>

Smith, A.M., S.B. Wilson, M. Thetford, K.L. Nolan, and C. Reinhardt Adams. 2014. "Performance of nine Florida native wildflower species grown in varying container substrates." *Native Plants*

Journal 15(1):75–86.

<https://doi.org/10.3368/npj.15.1.75>

Wilson, S.B., H. Perez, and M. Thetford. 2010. "Propagation, Production, and Landscape Evaluation of Native Wildflowers in West, Central and South Florida, 2010 Condensed Progress Report."

<https://www.flawildflowers.org/wp-content/resources/pdfs/Research/Wildflower%20year-end%20progress%20report-Wilson%20et%20al.pdf>

¹ This document is SGEB-75-4, one of a series of the Florida Sea Grant College Program, UF/IFAS Extension. Original publication date September 2018. Revised January 2025. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.

² Debbie Miller, professor of plant and wildlife community ecology, Department of Wildlife Ecology and Conservation, UF/IFAS West Florida Research and Education Center, Milton, FL; Mack Thetford, associate professor; Department of Horticultural Sciences, UF/IFAS West Florida Research and Education Center, Milton, FL; Christina Verlinde, former Extension agent II, Florida Sea Grant Program, UF/IFAS Extension Santa Rosa County, Milton, FL; Gabriel Campbell, graduate research assistant, UF/IFAS West Florida Research and Education Center, Milton, FL; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, dean for UF/IFAS Extension.