

Amelanchier x grandiflora 'Autumn Brilliance': 'Autumn Brilliance' Apple Serviceberry¹

Edward F. Gilman, Dennis G. Watson, Ryan W. Klein, and Deborah R. Hilbert²

Introduction

Apple serviceberry is a hybrid between *Amelanchier canadensis* and *Amelanchier laevis* that grows 15 to 25 feet tall. Its multiple stems are upright and highly branched, forming a dense shrub, or if properly pruned in the nursery, a small tree. It is superior to the species in that it suckers less and is adapted to a wide range of soils, but tolerates some drought. This cultivar supposedly resists leaf spot better than the original hybrid. The main ornamental feature is the spectacular white flowers that are larger than those of other amelanchiers. The flowers are borne in early spring and are at first tinged with pink but later fade to white. The young leaves are purplish and the fall color is red, yellow, or orange. Its edible fruit attracts birds. It is well adapted for planting along residential streets where there is plenty of soil space beneath power lines.

General Information

Scientific name: *Amelanchier x grandiflora*

Pronunciation: am-meh-LANG-kee-er x gran-dih-FLOR-uh

Common name(s): 'Autumn Brilliance' apple serviceberry

Family: *Rosaceae*




Figure 1. Middle-aged *Amelanchier x grandiflora* 'Autumn Brilliance': 'Autumn Brilliance' Apple Serviceberry

USDA hardiness zones: 3B through 7B (Figure 2)

Origin: not native to North America

Invasive potential: not assessed/incomplete assessment

1. This document is ENH237, one of a series of the Department of Environmental Horticulture, UF/IFAS Extension. Original publication date November 1993. Revised November 2023. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. Edward F. Gilman, professor emeritus; Dennis G. Watson, former associate professor, Department of Agricultural and Biological Engineering; Ryan W. Klein, assistant professor, arboriculture; and Deborah R. Hilbert, Gulf Coast Research and Education Center; Department of Environmental Horticulture; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, dean for UF/IFAS Extension.

Uses: container or planter; highway median; street without sidewalk; deck or patio; specimen; tree lawn 3–4 feet wide; tree lawn 4–6 feet wide; tree lawn > 6 ft. wide




Figure 2. Range

Description

Height: 15 to 25 feet

Spread: 15 to 25 feet

Crown uniformity: symmetrical

Crown shape: upright/erect, vase

Crown density: moderate

Growth rate: slow

Texture: fine

Foliage

Leaf arrangement: alternate (Figure 3)




Figure 3. Foliage

Leaf type: simple

Leaf margin: serrate

Leaf shape: elliptic (oval), oblong

Leaf venation: pinnate

Leaf type and persistence: deciduous

Leaf blade length: less than 2 inches, 2 to 4 inches

Leaf color: green

Fall color: red, orange

Fall characteristic: showy

Flower

Flower color: white/cream/gray

Flower characteristics: very showy

Fruit

Fruit shape: round

Fruit length: less than 0.5 inch

Fruit covering: fleshy

Fruit color: red

Fruit characteristics: attracts birds; showy; fruit/leaves not a litter problem

Trunk and Branches

Trunk/bark/branches: branches don't droop; showy; typically multi-trunked; thorns

Pruning requirement: little required

Breakage: resistant

Current year twig color: brown

Current year twig thickness: thin

Wood specific gravity: unknown

Culture

Light requirement: full sun, partial sun or partial shade

Soil tolerances: clay; sand; loam; acidic; well-drained

Drought tolerance: moderate

Aerosol salt tolerance: moderate

Other

Roots: not a problem

Winter interest: no

Outstanding tree: yes

Ozone sensitivity: unknown

Verticillium wilt susceptibility: unknown

Pest resistance: sensitive to pests/diseases

Use and Management

'Robin Hill' has an upright, tree-form growth habit, 20 to 25 feet tall, but is sensitive to drought.

Pests

Cambium miners cause concern when noticed but are not very damaging to the tree. The mines can extend from a twig all the way down to the roots. The mines are light colored lines in the bark. No controls are suggested.

A leaf miner will mine leaves, particularly the lower half of the leaf. The mines are irregular in shape.

The leaves of amelanchier are skeletonized by at least two insects. The first insect forms small cocoons on the undersides of leaves. Skeletonized leaves look as though they have windows in them after the insects scrape tissue off the top and bottom of the leaves. The second insect is the larva of the pear sawfly. The larvae are black to greenish black and look slimy. Adult sawflies lay eggs in May and June and again in August. Heavily skeletonized leaves drop off.

Several borers attack amelanchier. Healthy trees are considered less susceptible, so regular fertilization and watering during dry spells will help prevent borer attacks.

Spider mites will feed on amelanchier. These insects are hard to detect as they are so small. The main symptom of mite injury is the loss of green leaf coloration. If the infestation is heavy, very fine webbing may be seen. Horticultural oil sprays help control mite infestations.

Aphids of several types suck juices from amelanchier. Heavy infestations cause distortion of the foliage and new growth, and deposit large amounts of sticky honeydew on lower foliage. Black sooty mold will grow on the honeydew.

Diseases

Witches' broom, also called black mildew, infects the growing point causing the formation of many stems. The cluster of stems is called the "witches' broom". Another symptom is a black fungal growth, coating the undersides of the leaves. The damage to the tree is usually not serious and the brooms can be pruned off. No chemical controls are suggested.

Leaf blight can cause leaf drop when a severe infection occurs. The disease causes small purple spots on the leaves. The spots enlarge and turn brown, later a small black dot will be seen in the center of the spot. Large numbers of spots cause infected leaves to drop.

Fire blight is characterized by the sudden wilting and death of branch tips. The blossoms wilt, blacken, and hang on the twig. The bark is shriveled and has small bumps or blisters on it. Sometimes gum oozes out of the infected area and a crack forms between the diseased and healthy bark. Control with chemicals is difficult. Diseased branches should be pruned out. Make the cut at least four inches beyond the diseased area. Disinfect pruning tools with bleach between cuts. Fertilizing heavily with nitrogen increases susceptibility to fire blight.

Powdery mildews of several types cause white powdery growth on the leaves of amelanchier. Late in the season no controls may be needed.

Fruit rot may be a problem in wet weather. The fruits are often eaten by birds so may not be around long enough to become diseased.

Cedar rusts can be troublesome.