UF IFAS Extension UNIVERSITY of FLORIDA

ENH257

Betula pendula 'Youngii': 'Youngii' European Birch[1]

Edward F. Gilman and Dennis G. Watson[2]

Introduction

Young's weeping European birch is graceful, weeping ornamental with wonderful yellow fall color but is susceptible to fatal attacks of bronze birch borer. The leaves are often browned by birch leaf miner. When grown, plan to provide the necessary insect control and provide the necessary cultural conditions for best growth. It grows moderately with a variable height, depending on how it was trained. Some specimens appear sloppy, others more tailored. Lawn grasses grow fairly well in its light shade. A moist soil and a regular fertilization program plus watering in dry weather are suggested. Although popular, the tree requires more care and spraying than other ornamental trees. Not a low-maintenance tree. Maintaining a good mulch around the root zone is helpful for best growth.

General Information

Scientific name: *Betula pendula*
Pronunciation: BET-yoo-luh PEND-yoo-luh
Common name(s): 'Youngii' European birch
Family: *Betulaceae*
USDA hardiness zones: 3A through 6B (Fig. 2)
Origin: not native to North America
Invasive potential: little invasive potential
Uses: deck or patio; specimen; shade
Availability: not native to North America

Description

Height: 12 to 20 feet
Spread: 15 to 25 feet
Crown uniformity: irregular
Crown shape: weeping
Crown density: moderate
Growth rate: slow
Texture: fine




Figure 1. Middle-aged *Betula pendula* 'Youngii': 'Youngii' European Birch

Foliage

Leaf arrangement: alternate (Fig. 3)
Leaf type: simple
Leaf margin: double serrate
Leaf shape: rhomboid, ovate
Leaf venation: pinnate

1. This document is ENH257, one of a series of the Environmental Horticulture, UF/IFAS Extension. Original publication date November 1993. Revised December 2006. Reviewed February 2014. Visit the EDIS website at http://edis.ifas.ufl.edu.

2. Edward F. Gilman, professor, Environmental Horticulture Department; Dennis G. Watson, former associate professor, Agricultural Engineering Department, UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

Leaf type and persistence: deciduous
Leaf blade length: less than 2 inches, 2 to 4 inches
Leaf color: green
Fall color: yellow
Fall characteristic: showy




Figure 2. Range




Figure 3. Foliage

Flower

Flower color: brown
Flower characteristics: showy

Fruit

Fruit shape: elongated
Fruit length: .5 to 1 inch, 1 to 3 inches
Fruit covering: dry or hard
Fruit color: brown
Fruit characteristics: attracts birds; not showy; fruit/leaves not a litter problem

Trunk and Branches

Trunk/bark/branches: branches droop; not showy; typically multi-trunked; thorns
Pruning requirement: needed for strong structure
Breakage: resistant
Current year twig color: brown
Current year twig thickness: thin
Wood specific gravity: unknown

Culture

Light requirement: full sun
Soil tolerances: sand; loam; clay; acidic; well-drained
Drought tolerance: moderate
Aerosol salt tolerance: moderate

Other

Roots: not a problem
Winter interest: yes
Outstanding tree: no
Ozone sensitivity: unknown
Verticillium wilt susceptibility: resistant
Pest resistance: sensitive to pests/diseases

Use and Management

Several cultivars are listed, but these too will have pest problems: 'Dalecarlica'—deeply lobed leaves on pendulous branches; 'Laciniata'—cut leaves; 'Fastigiata'—upright growth habit; 'Purple Splendor' (purpurea), 'Scarlet Glory'—purple leaves; 'Tristis'—weeping habit; 'Youngii'—weeping habit.

Pests

A light aphid infestation may not be serious, but heavy infestations cause distorted and stunted growth and produce large amounts of honeydew. The honeydew serves as a substrate for sooty mold.

Birch skeletonizer feeding causes leaf browning. The skeletonizer larva is yellowish-green and one quarter-inch long.

Birch leaf miner is a common insect pest of birch. A small white worm eats out the middle of the leaf, which turns brown. Severe attacks of birch leaf miner predispose trees to bronze birch borer infestation. The insect shows up in mid-May, but timing can vary from one year to the next and will vary according to your location in the country. The first of two generations per year is the most damaging.

The most serious pest of landscape white birches is bronze birch borer. Stressed trees are most susceptible to borer attacks. The insect bores in the sapwood, beginning in the top third of the tree, causing death of the tree crown. The tunnels are slightly raised and faintly rust colored. Emergence holes in the trunk are shaped like capital Ds. Keep the trees healthy by controlling other insects, fertilizing, and watering as needed. Chemical control is applied to the trunk and main branches. Timing of the first spray will vary from year to year depending on weather conditions.

A commercial sprayer may be needed to apply the spray adequately.

Diseases

Several fungi cause canker diseases on birch. These diseases infect and kill sapwood, causing sunken areas on the trunk and larger branches. There is no chemical control for canker diseases. Preventive measures include keeping the tree healthy and avoiding wounding. Regular fertilization will keep birches vigorous and more resistant to cankers. Water in dry weather to prevent water stress.

Dieback is characterized by a slow death of the branches. The tree crown accumulates dead branches. Injury caused by bronze birch borer is similar but far more prevalent. Prevent dieback by maintaining tree vigor with water and fertilizer. When the disease does occur, prune out dead branches and increase tree vigor.

Several fungi also cause leaf spots that, when severe, can cause defoliation.