UF IFAS Extension UNIVERSITY of FLORIDA

ENH346

Cornus florida 'First Lady': 'First Lady' Flowering Dogwood[1]

Edward F. Gilman and Dennis G. Watson[2]

Introduction

The state tree of Virginia, flowering dogwood grows 20 to 35 feet tall and spreads 25 to 30 feet. This cultivar has tri-colored leaves with yellow and light green markings. Leaves turn mostly green during the summer in the southern part of its range. The tree can be trained with one central trunk or as a picturesque multi-trunked tree. The flowers consist of four bracts which subtend the small head of yellow flowers. The bracts are white and have a fall color of red to pink. The bright red fruits are often eaten by birds. Fall color is more vivid in USDA hardiness zones 5 to 8a. Branches on the lower half of the crown grow horizontally, those in the upper half are more upright. In time, this can lend a strikingly horizontal impact to the landscape, particularly if some branches are thinned to open up the crown. Lower branches left on the trunk will droop to the ground, creating a wonderful landscape feature.

General Information

Scientific name: *Cornus florida*
Pronunciation: KOR-nus FLOR-ih-duh
Common name(s): 'First Lady' flowering dogwood
Family: *Cornaceae*
USDA hardiness zones: 6A through 9A (Fig. 2)
Origin: native to North America
Invasive potential: little invasive potential
Uses: deck or patio; screen; specimen; shade; tree lawn 3–4 feet wide; tree lawn 4–6 feet wide; tree lawn > 6 ft wide
Availability: somewhat available, may have to go out of the region to find the tree



Figure 1. Middle-aged *Cornus florida* 'First Lady': 'First Lady' flowering dogwood
Credits: Ed Gilman, UF/IFAS

1. This document is ENH346, one of a series of the Environmental Horticulture Department, UF/IFAS Extension. Original publication date November 1993. Reviewed June 2014. Visit the EDIS website at http://edis.ifas.ufl.edu.

2. Edward F. Gilman, professor, Environmental Horticulture Department; Dennis G. Watson, former associate professor, Agricultural Engineering Department, UF/IFAS Extension, Gainesville FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.



Figure 2. Range

Description

Height: 20 to 30 feet
Spread: 25 to 30 feet
Crown uniformity: symmetrical
Crown shape: round
Crown density: moderate
Growth rate: slow
Texture: medium

Foliage

Leaf arrangement: opposite/subopposite (Fig. 3)
Leaf type: simple
Leaf margin: entire
Leaf shape: ovate
Leaf venation: pinnate, bowed
Leaf type and persistence: deciduous
Leaf blade length: 2 to 4 inches, 4 to 8 inches
Leaf color: variegated
Fall color: red
Fall characteristic: showy




Figure 3. Foliage

Flower

Flower color: white/cream/gray
Flower characteristics: very showy

Fruit

Fruit shape: oval, round
Fruit length: less than .5 inch
Fruit covering: fleshy
Fruit color: red
Fruit characteristics: attracts birds; showy; fruit/leaves not a litter problem




Figure 4. Flower



Figure 5. Fruit

Trunk and Branches

Trunk/bark/branches: branches droop; not showy; typically multi-trunked; thorns
Pruning requirement: little required
Breakage: resistant
Current year twig color: green
Current year twig thickness: medium
Wood specific gravity: unknown

Culture

Light requirement: full sun, partial sun, or partial shade, shade tolerant
Soil tolerances: sand; loam; clay; acidic; slightly alkaline; well-drained
Drought tolerance: moderate
Aerosol salt tolerance: low

Other

Roots: not a problem
Winter interest: no
Outstanding tree: no
Ozone sensitivity: unknown
Verticillium wilt susceptibility: resistant
Pest resistance: sensitive to pests/diseases

Use and Management

The tree is not suited for parking lot planting but can be grown in a wide street median, if provided with less than

full-day sun and irrigation. Dogwood is a standard tree in many gardens where it is used by the patio for light shade, in the shrub border to add spring and fall color or as a specimen in the lawn or groundcover bed. It can be grown in sun or shade but shaded trees will be less dense, grow more quickly and taller, and have poor fall color. Trees prefer part shade (preferably in the afternoon) in the southern end of its range. Many nurseries grow the trees in full sun, but they are irrigated regularly. Flowering dogwood prefers a deep, rich, well-drained, sandy or clay soil and has a moderately long life. It is not recommended in the New Orleans area and other heavy, wet soils unless it is grown on a raised bed to keep roots on the dry side. The roots will rot in soils without adequate drainage.

Several of the cultivars listed are not readily available: Pink-flowering cultivars grow poorly in USDA hardiness zones 8 and 9. 'Apple Blossom'—pink bracts; 'Cherokee Chief'—red bracts; 'Cherokee Princess'—white bracts; 'Cloud 9'—white bracts, many blooms, flowers at early age; 'Fastigiata'—upright growth while young, spreading with age; 'Gigantea'—bracts 6-inches from tip of one bract to tip of opposite bract; 'Magnifica'—bracts rounded, 4-inch diameter pairs of bracts; 'Multibracteata'—double flowers; 'New Hampshire'—flower buds cold hardy; 'Pendula'—weeping or drooping branches; 'Plena'—double flowers; var. *rubra*—pink bracts; 'Spring Song'—bracts rose red; 'Springtime'—bracts white, large, blooms at an early age; 'Sunset'—supposedly resistant to anthracnose; 'Sweetwater Red'—bracts red; 'Weaver's White'—large white flowers, adapted to the south; 'Welchii'—leaves variegated with yellow and red; 'White Cloud'—flowers more numerous, bracts white; 'Xanthocarpa'—fruit yellow.

Pests

Aphids may be controlled by spraying them with a strong stream of water from the garden hose.

Several borers will attack dogwood. Try to keep the trees healthy with regular fertilization, and irrigation during dry weather. Indications of borer problems are holes in the trunk, leaves smaller than normal, and dieback of the crown.

Dogwood club gall midge causes galls at the branch tips. The leaves on affected branch tips may be distorted and the branch may fail to form a flower bud. Prune out the galls as soon as they are seen.

Leaf miners cause brown blister-like mines on the undersides of leaves. The adult leaf miner skeletonizes the leaves. Scales can build up to large numbers before being detected. Horticultural oil will help control overwintering stages.

Twig girdlers prune the tips of small branches. They are more of an annoyance than a serious problem unless you are a nursery operator.

Diseases

Dogwood anthracnose may be the biggest concern with growing flowering dogwood. Infection is favored by cool, wet spring or fall weather. Drought and stressed trees appear to be most affected, as are those at higher elevations. Trees on the coastal plain may be much less likely to become affected. Consecutive years of infection can kill trees. Keep the tree healthy with regular irrigation in dry weather but avoid overhead irrigation. Plant the tree in an area which allows the leaves to stay as dry as possible. Consult a local pathologist for the latest in control measures. *Cornus kousa* is thought to be resistant to anthracnose and it can be planted in areas where anthracnose is a problem. It is a very beautiful tree.

Early symptoms of dogwood canker are smaller and paler leaves than normal. Leaves on infected branches are red earlier in the fall. At first the symptoms appear only on the infected side of the tree but become more general as the canker enlarges. There is no chemical control for the disease. Avoid trunk wounds during and after planting.

Crown canker is associated with wet soils and can be controlled with appropriate fungicides.

Flower and leaf blight attacks fading bracts, especially during wet weather. Infected flower parts fall on the leaves spreading the infection.

A large number of leaf spots attack dogwood. Clean up and dispose of infected leaves.

Powdery mildew covers the leaves with a fine white coating.

Leaf scorch occurs during hot, dry, windy weather. This condition looks like a disease. Scorch symptoms are drying and browning of the leaf margins, or, in more serious cases, drying and browning of the interveinal area.